

THE REPUBLIC OF SUDAN  
FOREST NATIONAL CORPORATION

FOREST REFERENCE LEVEL SUBMISSION TO THE  
UNFCCC



Food and Agriculture  
Organization of the  
United Nations



GREEN  
CLIMATE  
FUND

June 2025



## FOREWARD

The Republic of Sudan (RoS) is one of the most vulnerable countries to climate change and its devastating impacts and therefore, highly committed to the international cooperation on addressing this phenomenon. REDD+ is among the top priority programmes for RoS to participate in the global mitigation efforts and to build resilience in its land sector. RoS is integrating the global REDD+ objectives into its national climate change related plans and strategies.

The RoS NDC (2021) aims towards implementing low carbon development interventions in three sectors - energy, forestry and waste. Sustainable management of degraded forest and the Gum Arabic belt; afforestation and restoration of degraded agriculture lands; and restoration/conservation of mangrove forests along Red Sea coast; are among the key contributions included in the NDC (2021).

In 2021 RoS adopted a National REDD+ Strategy (NRS), with the objective of maximizing carbon and non-carbon benefits through improved sustainable natural resources management. The NRS support efforts to achieve REDD+ objective and contribute to the global climate change mitigation in the context of sustainable development. In this context, the preparation of this forest reference level (FRL) will support and enable effective implementation of the national REDD+ Strategy (NRS), RoS's NDCs, monitoring and reporting on the forest sector mitigation actions.

The submission of this FRL is a clear demonstration of RoS commitment and keen interest to undertake ambition climate mitigation actions and transition its forest sector towards a low emission, sustainable development pathway.



Anwar Abdelhameid  
Director General, Forest National Corporation

## ACKNOWLEDGEMENT

The preparation of this National Forest Reference Level (FRL) of Republic of Sudan (RoS) under the current war circumstances demonstrate a high level of determination, dedication and commitment of Sudanese Government institutions and experts to pursue their international obligations.

The successful completion of this submission is a result of collaboration and contribution of national institutions and experts, without their dedication it would have been impossible to achieve this success.

I would like to acknowledge with much gratitude the crucial role of the National REDD+ Steering Committee for the wise guidance they provided throughout the preparation of the national FRL.

My sincere thanks and gratitude to the members of the FRL Taskforce and Remote Sensing Team for their valuable contributions, dedication and commitment to successfully achieving all their assigned tasks and activities despite the difficulties in communications, internet connection and access to offices.

I am greatly indebted to the national and international consultants who provided the needed technical support and backstopping to the various stages of the FRL preparation.

I am extremely grateful to Mr Anwar Abdelhameid, the Director General of the Forest National Corporation (FNC) who provided full support and guidance which enable the REDD PMU to successfully complete this FRL submission despite the challenging war circumstances.

I would like to present my warmest thanks and gratitude to all colleagues of FNC staff, specially colleagues in the different States of the country who provided expert input and supported the data collection phase of this FRL preparation, they really deserved special thanks and appreciation.

Last but not least, I would like to express my sincere gratitude and appreciation to colleagues of FAO RoS and Rome, for their continuous support, valuable input and guidance that enabled us to timely complete this FRL submission.



Dr. Sayeda Ahmed Khalil  
Coordinator, REDD+ Programme Management Unit

## ACRONYMS

AD	Activity Data
A/R	Afforestation and reforestation
AGB	Above-ground biomass
ABGD	Above-ground biomass density
BGB	Below-ground biomass
BUR	Biennial Update Report
CEO	Collect Earth Online
CO <sub>2</sub>	Carbon Dioxide
CO <sub>2</sub> e	Carbon Dioxide Equivalent
GEDI	Global Ecosystem Dynamic Investigation
FAO	Food and Agriculture Organization of United Nation
FNC	Forests National Corporation
FREL	Forest Reference Emission Level
FRL	Forest Reference Level
GDP	Gross domestic product
GHGs	Green House Gases
GLCN	Global land cover network
Ha	Hectare
HCENR	Higher Council for Environment and Natural Resources
NDC	National Determined Contribution
IPCC	Intergovernmental Panel on Climate Change
LULUCF	Land use land use change and forestry
NDCs	National determined contribution
NFI	National Forest Inventory
NFMs	National Forest Monitoring System
NWFP	Non woody forest products
REDD+	Reducing Emissions from Deforestation and forest Degradation; and the role of conservation, sustainable management of forests, and enhancement of forest carbon stocks
RSSA	Remote Sensing and Seismology Authority
SLC	Scan Lune Corrector
SNC	Second National Communication
SU	Sampling Unit
t.CO <sub>2</sub>	Tonnes of Carbon Dioxide
TNC	Third National Communication
UNFCCC	United Nations Framework Convention on Climate Change

## TABLE OF CONTENTS

<b>FOREWARD</b>	<b>2</b>
<b>ACKNOWLEDGEMENT</b>	<b>3</b>
<b>ACRONYMS</b>	<b>4</b>
<b>TABLE OF CONTENTS</b>	<b>5</b>
<b>LIST OF TABLES</b>	<b>7</b>
<b>LIST OF FIGURES</b>	<b>8</b>
<b>SUMMARY</b>	<b>9</b>
<b>1. INTRODUCTION</b>	<b>11</b>
<b>2. FOREST DEFINITION</b>	<b>15</b>
<b>3. CONSISTENCY WITH GHGs INVENTORY REPORTING</b>	<b>15</b>
<b>4. SCALE OF THE FRL</b>	<b>16</b>
4.1. The Historical Reference Period (RP):	16
<b>5. SCOPE OF THE FRL</b>	<b>18</b>
5.1. REDD+ Activities of the FRL:	18
5.2. Carbon pools:	20
5.3. Greenhouse Gases:	21
<b>6. ACTIVITY DATA FOR DEFORESTATION</b>	<b>22</b>
6.1. Sampling Design	23
6.1.1 Systematic NFI grid based on permanent stratification	23
6.1.2 Intensification of the sampling grid	25
6.1.2.1 Training data collection	25
6.1.2.2 Provision of satellite image composites	26
6.1.2.3 Forest Non-Forest (FNF) Probability Maps	27
6.1.2.4 Change probability Layer	29
6.1.2.5 Stratification of change probability	29
6.1.2.6 Sample Allocation	29
6.3. Data Collection	30
6.4. Capacity Building	33
6.5. Results	33
6.5.2. Results by States	35
6.6. Quality Management	39
<b>7. Description of the National Forest Inventory (NFI):</b>	<b>41</b>
7.1. NFI Sampling Design:	41
7.2 Data Collection Process	42
7.3. Data Analysis	43
<b>8. Emission Factors for Deforestation:</b>	<b>44</b>
8.1. Revised estimates of above-ground biomass:	44
8.1.2. Methodology	45
8.1.3. Results	52
8.2. Dead wood:	56

8.3. Emission Factors for deforestation:	56
<b>9. Emission Factors for Forest Degradation:</b>	<b>60</b>
<b>10. The Activity Data for Enhancement of Forest Carbon Stock:</b>	<b>63</b>
10.1. Removal Factors for Enhancement of Forest Carbon Stock:	66
<b>11. APPROACHES AND METHODS APPLIED IN THE FRL CONSTRUCTION</b>	<b>70</b>
11.1. Estimation of GHGs Emission from Deforestation:	70
11.1.1. Uncertainty:	72
11.2. Estimation of GHGs Emission from Degradation:	72
11.3. Estimation of Removal from Enhancement of Forest Carbon Stock:	73
<b>12. RELEVANT POLICIES AND LEGISLATION</b>	<b>76</b>
<b>13. PROPOSED FOREST REFERENCE LEVEL</b>	<b>78</b>
13.1. Values of national FRL:	78
13.2. Description of change from Sudan's FRL 2020:	80
<b>14. ADJUSTMENT FOR NATIONAL CIRCUMSTANCES</b>	<b>82</b>
<b>15. UPDATING FREQUENCY</b>	<b>82</b>
<b>16. FUTURE IMPROVEMENTS</b>	<b>82</b>
<b>17. REFERENCES</b>	<b>85</b>
<b>18. ANNEXES</b>	<b>88</b>
<i>Annex 1: Areas of the Afforestation and Reforestation of the 18 States of RoS</i>	<b>88</b>
<i>Annex 2: Wood Density data</i>	<b>90</b>
<i>Annex 3: Sampling unit, plot and subplot design</i>	<b>94</b>
<i>Annex 4: Calculation Procedures Used in the RoS NFI</i>	<b>95</b>
<i>Annex 5: List of Assets used for Activity Data</i>	<b>102</b>
<i>Annex 6: Annual Composite of Landsat Year 2011</i>	<b>103</b>
<i>Annex 7: A visual of the “stable forest”, “forest degradation” or “forest gain” locations of the NFI</i>	<b>105</b>
<i>Annex 8: Response Design</i>	<b>118</b>
<i>Annex 9: Members of Steering Committee, FRL Taskforce, Remote Sensing Team and Management and Technical Support Team</i>	<b>125</b>

## LIST OF TABLES

Table 1: Republic of Sudan's (RoS) FRL compliance with the relevant UNFCCC decisions.....	9
Table 2: The Strategic Options and PAMs of the NRS .....	14
Table 3: The carbon pools included and omitted in the national FRL.....	21
Table 4: Description of the Strata, along with total area covered by each stratum (Source: NFI Report Table 3.2) .....	23
Table 5: NFI Sampling Design: Distribution of Sample by Stratum and total filed visited samples during NFI cycle 2017 -2020. (Source: NFI Report Table 3.2).....	24
Table 6: Categorization of training data into Forest and non-Forest class .....	26
Table 7: Depicting details of Optical and Radar composite used for classification along with relevant bands. .....	28
Table 8: Sample intensification by stratum along with total area of stratum in hectares .....	30
Table 9: Stratified Area Estimates by States in Hectares .....	36
Table 10: Annual Areas of Deforestation by State in Hectares .....	37
Table 11: Annual Areas of Forest Degradation by State in Hectares .....	38
Table 12: Disaggregation of deforestation area by LULC classes .....	39
Table 13: Matrix of by LULC classes with majority agreement .....	40
Table 14: Average above-ground biomass density by state.....	55
Table 15: Average aboveground volume/hectare (v/ha) of Standing deadwood disaggregated by state .....	56
Table 16: Weighted Average Wood Density Values of the main Tree Species per State .....	57
Table 17: Emission factors for deforestation per State .....	59
Table 18: Average percentage reduction in canopy cover of forest degradation area per State... ..	61
Table 19: Emission factors for forest degradation per State.....	62
Table 20: Success rates of A/R forests planted by Seeds and Seedlings for each State .....	65
Table 21: Common tree Species used in the annual A/R activities and percentages of their areas in each State .....	66
Table 22: Mean Annual Increment values and other parameters used in estimation of A/R removals.....	67
Table 23: Removal Factors, Weighted Average Annual Biomass Carbon Increment .....	69
Table 24: Proposed Values of the Deforestation FREL.....	78
Table 25: Proposed Values of the Degradation FREL.....	79
Table 26: Proposed Values of the Enhancement of Forest Carbon Stock FRL.....	79
Table 27: Comparison of Sudan FRL 2020 and FRL 2025 .....	80

## LIST OF FIGURES

Figure 1: Republic of Sudan Map .....	12
Figure 2: Geographic distribution of sample intensities over each stratum .....	24
Figure 3: Showing workflow of the activity data .....	28
Figure 4: Customized Response Design established for Republic of Sudan for data collection ..	32
Figure 5: Collect Earth Survey Card used for activity data collection using Collect Earth software .....	33
Figure 6: Annual deforestation trend from 2011-2023 .....	35
Figure 7: Showing trend of forest changes by states over 2012 -2021 .....	37
Figure 8: <i>Showing comparison of top five interpreter average with overall average by LULC and forest change classes. Table 1 refers to LULC overall average by classes, Table 2- is showing average of top five interpreter by LULC classes. Table 3- is showing forest change classes overall average. Table 4 – showing an average of first five interpreters by forest change classes.</i> .....	40
Figure 9: Land use Land Cover Section .....	42

## SUMMARY

This document presents Republic of Sudan's (RoS) National Forest Reference Level (FRL), which is submitted for the technical assessment under the UNFCCC. Table 1, below provides summary description of the FRL submission and its consistency with the relevant UNFCCC guidance and summarizes the decisions made by the government of Sudan in relation to the scale, scope and methods used in FRL construction. This FRL represents forest conditions in the dry lands of Sub Sahara Africa. The FRL includes three REDD+ activities of reducing emissions from deforestation, reducing emissions from forest degradation and Enhancement of Forest Carbon Stock. Deforestation occurs in 13 states, with national average annual emission associated with deforestation estimated at **9,317,358 tonne CO<sub>2</sub> /Year**. Forest degradation occurs in 12 states, with national average annual emission associated with forest degradation estimated at **1,678,433 tonne CO<sub>2</sub>/Year**. The enhancement of forest carbon stock covers 18 states, resulting in national average annual CO<sub>2</sub> removals of about - **2,112,017 tonne CO<sub>2</sub>/Year** due to Afforestation and Reforestation activities. The reference period for the three REDD+ activities in 2012-2021.

**Table 1: Republic of Sudan's (RoS) FRL compliance with the relevant UNFCCC decisions**

UNFCCC reference	Description	National FRL (2024)
Decision 12/CP.17 Paragraph II.10	Stepwise approach	National Scale disaggregated by States Builds on lessons learnt, data, methods and capacity developed by the FRL 2020 Reference historical period is 2012-2021
Decision 12/CP.17 Annex, paragraph (c)	Pools and gases	Pools: Aboveground, below ground biomass and standing deadwood Gases: CO <sub>2</sub>
Decision 12/CP.17 Annex, paragraph (c)	REDD Activities	<ul style="list-style-type: none"><li>Reducing emissions from deforestation</li><li>Enhancement of forest carbon stocks</li><li>Reducing emissions from forest degradation</li></ul>
Decision 12/CP.17 Annex, paragraph (d)	Forest definition and consistency with GHG inventories	Forest means an area of land spanning at least a minimum area of 0.4 ha with trees that have attained, or have the potential to attain at least 2 m. in height and a minimum tree canopy cover of 10%. It includes wind-breaks and/or shelter-belts with a minimum of 20 m. in width. The same definition used in GHGs inventory of the Third National Communication


Decision 12/CP.17 Annex	Guidelines and Methodologies	IPCC 2006 Guidelines for national GHGs inventories and data 19R IPCC guidelines The same methodology used in GHGs inventory of the Third National Communication
Decision 12/CP.17, Paragraph II.9	Information on rationale, national circumstances and their consideration in the national FRLs	Description of national circumstances provided No adjustment is needed, The reference period is considered representatives of the current trend and capturing key milestone in national circumstances affecting forest resources in the country

## 1. INTRODUCTION

Republic of Sudan (RoS) is submitting this national Forest Reference Level (FRL) in response to the invitation of the Conference of Parties to the UNFCCC for developing countries to develop and submit, on a voluntary basis, FREL/FRL, for consideration by the UNFCCC. Following the stepwise approach, stipulated in decision 12/CP.17, para II.10, RoS prepared this National scale Forest Reference Level (FRL), building on the experiences, capacities, data and resources developed during the preparation of its subnational FRL (2020) as well as the feedback received through the UNFCCC technical assessment. This submission is intended for technical assessment in the context of results-based payments for reducing emissions from deforestation and forest degradation and the role of conservation, sustainable management of forests and enhancement of forest carbon stocks in developing countries (REDD+) under UNFCCC (Para13 decision 12/CP.16 and para 71(b) decision 1/CP.16).

The objective of preparing the national Forest Reference Level (FRL) is to support achieving the overall objectives of the national forest programme and enhancing the contribution of the forest resources to global climate change mitigation in the context of sustainable development. This is to be achieved through enabling effective implementation of the national REDD+ Strategy (NRS) and national determined contribution (NDC) under the Paris Agreement. In its NDC (2021) RoS plans to implement low carbon development interventions in three sectors - energy, forestry and waste. The measures envisioned to achieve GHG emission reductions are consistent with RoS's national development priorities, objectives and circumstances. These measures include implementation of the National REDD+ Strategy including restoration and sustainable management of degraded forest reserve and the Gum Arabic belt; Afforestation and restoration of degraded lands including 10% of rainfed and 5% of irrigated areas of the agriculture land; Restoration/conservation of mangrove forests along Red Sea coast.

Located in North Eastern Africa, the RoS is bound by Egypt, The Red Sea, Eritrea, Ethiopia, Republic of South RoS (RSS), Central African Republic, Chad and Libya, with an estimated total of 1.882 million km<sup>2</sup> (Figure1). It lies between latitudes 100 and 220 N and longitudes 220 to 380 E. The highest point in the country is Jebel Marra; 3,024 meters above sea level (m.a.s.l.) and the lowest is the Red Sea. The most salient geographical features are the Nubian and Bayuda Deserts in the north, the Nile Valley, Jebel Marra, Nuba, and Ingessena & Red Sea Hills. The Blue Nile and White Nile Rivers originates in the Ethiopian Highlands and the Equatorial Lakes, respectively. The two rivers unite at Khartoum and form the River Nile that runs north to the Mediterranean Sea. Administratively RoS is divided in 18 State, see map in figure (1) below:



**Figure 1: Republic of Sudan Map**

The ecological zones of RoS extend over a wide range from the desert in the extreme north to the high rainfall savannah in the south. The country is ecologically divided into five vegetation zones following the rainfall patterns from North to South:

1. Desert: (0-75 mm of precipitation),
2. Semi-desert: (75-300 mm),
3. Low rainfall savannah on clay and sand: (300- 800 mm),
4. High rainfall savannah (800-1500 mm),
5. Mountain Vegetation: (300-1000 mm).

The country is rich in biodiversity with diverse environmental systems making it endowed with flora and fauna which are being subjected to a number of threats as a result of natural factors and human activities. According to the Land Cover Atlas of RoS, FAO (2012), Forests together with Rangeland represent 35.6% of the total country area. Forests play an important social, economic and environmental role by the goods and services they provide to support people's livelihoods. Approximately 63% of the population is rural and considered forest dependent essentially for fuelwood as main source of energy and round timber as building material. In addition to wood

products, the rural population uses the forest extensively for grazing, as source of bush meat (wildlife hunting) and for food in the form of tree leaves, fruits, honey and tubers. They also use tree shade for their social functions and other recreational purposes, (FOSA, 2001). Many other non-wood forest products are collected from forests for domestic as well for commercial purposes, among these, RoS is the leader country in gum Arabic production, other incenses, tannin, etc. Forest based small industries constitute a main pillar to the national economy as well as to local economies, and provide 15% of job opportunities in the country. In addition, more than five million farmers earn their living depending on Gum Arabic production. Forests contribute 30% of the fodder requirements of the national animal herd and this contribution can reach 70% in extremely dry years.

However, deforestation remains key threat to forest resources, according to the report on Drivers of Deforestation and Forest Degradation (DoDD) (2018), deforestation has caused RoS's forest cover to decline from 76.4 million hectares (ha) in 1990 to 70.49 million ha in 2000 and 69.95 million ha in 2010 (30.5% to 28.1% and 27.9% of the country total area, respectively (FRA, 2010). RoS Cover Map report (2020) estimates forest area at 22.8 million hectares, about 12% of the total country area. The significant reduction estimated in the Land Cover Map report is also attributed to the cessation of Sudan in 2011 into two States, the Republic of Sudan (RoS) and South Sudan, where more than half of the forest area fall in South Sudan. The estimation of the Land Cover Map (2020) is based on the new forest definition, mentioned in section 2 below, which reduces the dimensions of the area, highest and crown cover.

The conversion of the forests to agriculture remains the key driver of deforestation in the country, the DoDD (2018), estimated that agriculture expansion causes 42.1%. Agriculture expansion on forestland is caused by declining productivity of cropland, poor cultivation practices and population growth (2.8% per year in 2020). Over drivers, unsustainable extraction of fuelwood and charcoal for energy causes 19.8% and animal overgrazing causes 15.1% of the deforestation and the forest degradation in RoS.

RoS is one of the most vulnerable countries to climate change and hence accords high priority to the international cooperation to fighting climate change and avoiding its devastating consequences. REDD+ is among the top priority programmes in RoS to participate in the global mitigation efforts and to build resilience in its land sector, which host about 70% of the population. RoS is committed to the objectives of the global REDD+ programme and is working towards integrating these into its national forest programme and management practices. The National REDD+ Strategy (NRS) (2021) vision is to achieve sustainable management for a green economy while contributing to RoS's prosperity, livelihoods, and wellbeing. This is to be achieved through building synergies

with, and contribute effectively to, the National Development Strategy, which aims towards achieving a green prosperous country by 2030, while maximizing carbon and non-carbon benefits through improved sustainable natural resources management. The Strategy defines the general direction that the RoS will take to achieve the vision of the REDD+ programme. The NRS outlines five broad strategic options (Table 2) to be implemented through a set of policies (PAMs) and actions that collectively address the direct and indirect drivers of forest deforestation and degradation.

**Table 2: The Strategic Options and PAMs of the NRS**

STRATEGY OPTION	
<b>STRATEGY OPTION 1: INTEGRATED FOREST LANDSCAPE MANAGEMENT</b>	
PAM 1	Strengthen sectoral policies, financing, and institutional capacity
PAM 2	Strategic landscape management, restoration and emission reductions
PAM 3	Support for forestry research and development
<b>STRATEGY OPTION 2: CLIMATE SMART AGRICULTURE AND RANGELAND MANAGEMENT</b>	
PAM 4	Improving the adaptive and climate mitigation capacity of the agriculture sector
PAM 5	Promoting sustainable livestock and rangeland management
<b>STRATEGY OPTION 3: INTEGRATED LAND USE PLANNING</b>	
PAM 6	Harmonizing land use planning, investment policies, and legislation
PAM 7	Sustainable land management stewardship through land tenure security
<b>STRATEGY OPTION 4: SUSTAINABLE ENERGY SUPPLY AND USE</b>	
PAM 8	Increasing access to efficient and sustainable household energy
PAM 9	Promoting a sustainable biomass-based energy value chain
<b>STRATEGY OPTION 5: PROMOTING PARTICIPATION IN CLIMATE CHANGE RESPONSES</b>	
PAM 10	Advance the participation of youth and women

## **2. FOREST DEFINITION**

RoS's defines forest as an area of land spanning at least a minimum area of 0.4 ha with trees that have attained, or have the potential to attain at least 2 m in height and a minimum tree canopy cover of 10%. It includes wind-breaks and/or shelter-belts with a minimum of 20 m in width". This forest definition has been developed after cessation of RoS in 2011 into two states of RoS and South RoS. The rationale behind the new definition is to enhance the protection and production functions of the remaining forest resources. The new definition also responds to climate change requirement for forest definitions and enables realization of the role the forest resources envisioned to play in meeting RoS's obligations under the UNFCCC and Paris Agreement.

The forest definition was also used in the last national forest inventory (NFI Report 2021) and in the GHG inventory prepared for RoS's Third National Communication (TNC) to the UNFCCC and RoS's first Biennial Update Report (BUR).

The RoS does not have official definitions for the five REDD+ activities. However, for the purpose of this submission it is agreed that, within the circumstances of RoS, deforestation occurs as a result of conversion of forestland to other land use, mainly to cropland. Afforestation occurs on land that was not forest before, such as cropland, abandoned, degraded lands, etc, while reforestation occurs on bare lands in degraded forests areas that was cleared of their tree cover through human activities, mainly cultivation, and the cleared areas cannot regenerate naturally without human intervention, because of various reasons including the continuation of the same activities that causes their clearance.

## **3. CONSISTENCY WITH GHGs INVENTORY REPORTING**

This FRL submission is consistent with the GHGs inventory reports prepared for RoS's Third National Communication (TNC) and first Biennial Update Report (BUR). In particular:

- TNC, BUR and this FRL submission apply the same IPCC 2006 guidelines
- The same national and international data sources (FNC and research) of Afforestation and reforestation, wood density and other parameters are also used in this FRL submission and TNC/BUR.
- NFI (2021) data has been used by the FRL submission and the first BUR submission
- The same Forest definition has been used in the FRL submission and in TNC/BUR submission

The publication of the TNC and first BUR was planned in 2021, however, because of number of internal reasons including the current unrest situation this has been delayed till April 2025.

## **4. SCALE OF THE FRL**

The conclusions of the UNFCCC technical assessment on RoS subnational FRL 2020, identified one technical area for future improvement relevant to the scale of RoS's next FRL, which is the need to consider inclusion of other forest biomes as part of the efforts to construct a national FREL or FRL;

Accordingly, RoS decided to prepare this FRL as a national scale level covering all forestland and all land areas potentially available for enhancement of forest carbon stock in country. The estimation of emission associated with deforestation REDD activity covers all the areas of forestland in the country meeting the national forest definition. The estimation of the removals associated with the REDD activity of enhancement of forest carbon stock covers all areas of Afforestation and Reforestation (A/R) meeting the national forest definition in the country.

Almost 98% of the forestlands are located in the southern part of the country, within strata 3 and 4, the remaining about 1% are mostly Riverine forests located in strata 5 which is the area along the River Nile and its tributaries intersecting all other strata from south to north. Strat 1 and 2 represent the desert and semi-desert part. See the figure 2 in section 6.1, which is a map that shows the distribution and location of the forestland in the different stratum:

The national FRL will be disaggregated by States (subnational administrative units), consistent with the Forest National Corporation institutional arrangements, REDD management arrangements and the NFMS/MRV system. This will facilitate implementation of the national REDD+ strategy, emission reduction programmes/projects and the engagement of stakeholders at subnational levels. The deforestation FREL covers all the States with forest loss (13 States). While the enhancement FRL covers all land potentially available for implementing afforestation and reforestation activities in the 18 states of Sudan.

### **4.1. The Historical Reference Period (RP):**

The UNFCCC requires historical data to be taken into account in the construction of FREL/FRLs, however, it does not prescribe the length or the period that countries should use in the analysis of historical data. The historical period should provide for a realistic and robust benchmark to assess the mitigation performance of REDD+ activities. Ideally, it should represent the development,

dynamics and current trends in historical emissions. Number of REDD+ supporting and funding agency prefers 10 years reference period representing the most recent trend in the development of the forest sector.

In selecting the RP for this national FRL, RoS considered selecting a timeframe that encompasses key historical events, drivers and policies resulting in major impacts on the development of its forest sector. In this regards the secession of South Sudan in 9 July 2011 is considered the most important historical event affected forest resources. It was roughly estimated that about 60% of forestland and one third of the population become part of the state of South Sudan. As a result, more than 60% of Sudan's population are currently dependent on the remining forest resources. Given the fact that the population of the RoS is highly dependent on biomass energy (more 60% of the energy used), agriculture and livestock husbandry (more than 70% of the population). This situation has led to high pressure on the remaining forest resources and increases the rates of deforestation and forest degradation.

The national REDD+ Readiness Programme has established NFMs/MRV arrangements, a sub-national FRL, a Safeguards information system and adopted a National REDD+ Strategy during the period 2017-2021. All these are also important milestones for REDD+ implementation and for consideration in reference period determination.

The current forest Act issued 2002, is also an important development in the forest management and protection, it provides more consideration towards an integrated management approach including other natural resources such as rangelands. The Forest Act has provisions on allocation of 10% of the rainfed and 5% of the irrigated agricultural schemes to forestry and tree planting, aiming towards reversing deforestation, addressing land degradation and the poor productivity of cropland area, particularly in the mechanized rainfed areas which were originally forested area cleared of their tree cover and were subjected to mal-cultivation practices and are currently mostly degraded or degrading lands.

Taking into account all above-mentioned considerations, the RoS decided to apply and a reference period covering the timeframe of 2012 – 2021, which is representative of the current trends in the forest sector, it takes into consideration the important historical development mentioned above as well as allowing the possibility to use recent data and methods becoming available since the last sub-national FRL submission.

## 5. SCOPE OF THE FRL

Following the UNFCCC stepwise approach, decision 12/CP.17, para II.10, the scope of RoS's subnational FRL 2020 has included the most significant REDD activities, pools and gases. The REDD activities of deforestation and enhancement of forest carbon stock, the above and below ground biomass carbon pools and Carbon Dioxide was the only gas included. The key limitations to expand this scope to cover other activities, pools and gases were lack of good quality data, resources, e.g. to assess carbon stock change due to forest degradation and the significance of the contribution of other pools (e.g. deadwood, soil, litter). The conclusions from the UNFCCC technical assessment found the omission of some activities, pools and gases in 2020 FRL justifiable, however, identified three technical areas of improvement relevant to the scope of the national FRL, these include:

- Inclusion of other REDD+ activities, in particular reducing emissions from forest degradation, when new, adequate data and better information become available;
- Treatment of emissions from the deadwood, litter and soil pools
- Treatment of emissions of non-CO<sub>2</sub> gases (to maintain consistency with the national GHG inventory)

### 5.1. REDD+ Activities of the FRL:

The two REDD+ activities of reducing emissions from deforestation and enhancement of forest carbon stock, remain the most important and significant REDD+ activities in RoS. There are reliable data, information and experience for estimating them with more accuracy and therefore are included in the national FRL.

Forestland conversion to cropland remains key driver of deforestation, it is estimated that agriculture expansion causes 42.1% of forest loss. Agriculture expansion on forestland is caused by declining productivity of cropland, poor cultivation practices and population growth (2.8% per year in 2020). The DoDD (2018), made some important conclusion based on an analysis of a time series data of 63 years (1953-2016) of the cultivated area, production and productivity of the major food commodities (cereals and oilseeds). The area cultivated increased from an average of 5.6 million feddans to 41.3 million feddans during this period, indicating an increase of more than seven folds. The percentage of the area harvested decreased from an average of 90% to only 69% during the same period, implying a loss of productivity of almost 31% of the area under cultivated. The production increased during this period by about four folds from 1.5 million metric tons 6.7

million metric tons. This result indicates clearly that the country is relying on horizontal area expansion for food security.

On the other hand, RoS has a significant potential to implement the REDD activity of enhancement of forest carbon stock, due to the size of its land area and the vast forestlands cleared for agriculture production during the time since 1940s-1990s and after, most of these areas are left degraded or are degrading now. The potential land available for afforestation and reforestation is estimated at more than 9 million ha, including bare lands inside degraded forest areas, degraded area within the Gum Arabic belt, 5% the area of the irrigated agriculture schemes and 10% the areas of the rainfed agriculture schemes, the latter two are based on the provision in the Forest Law (2002). The estimated land area of rainfed sector does not include the traditional rainfed sector, which has the largest share of about 60% of the total rainfed area in the country. Data on the areas of the annual afforestation and reforestation activities is available in the records and annual reports of FNC HQ and its offices in the 18 states of the country.

As stated in the 2020 FRL submission and discussed during the UNFCCC technical assessment, forest degradation is also a significant REDD activity in Sudan. There is no official national definition of forest degradation. However, the DoDD (2018) report described it as “a gradual process through which a forest's biomass declined, its species composition changed, or its soil quality declined”. According to the same report, forest degradation is mainly caused by unsustainable wood extraction and overgrazing. These two activities are significant in Sudan due to the high dependence on wood for energy and the large animal population. The activity data for forest degradation has been developed through remote sensing means as explained in section 6 below. Forest degradation has been identified in 12 states, Table 11 below shows the area of forest degradation by state.

However, the key issue Sudan faces in the estimation of emissions associated with forest degradation is the lack of suitable data to develop emission factors. RoS does not have historical data and records from regular NFI measurements. Based on the discussion during technical exchange of UNFCCC technical assessment, Sudan decided to include forest degradation activity in this submission. The EFs for estimating emissions from forest degradation is based on the percentage reduction of forest canopy cover of the area affected by forest degradation, estimated during the reference period. Further description on the methodology for estimating emission associated with forest degradation is provided in section 9 below.

Regarding the inclusion of the other REDD+ activities of sustainable management of forests and conservation of forest carbon stock. Currently there is no data to assess emissions/removals associated with these activities in RoS.

## **5.2. Carbon pools:**

The UNFCCC decision 12/CP.17, Annex paragraph C, requires significant carbon pools to be included in the FREL/FRLs, otherwise reasons of their omission should be provided. In its 2020 FRL submission, RoS included above and below ground biomass carbon pools and provided the reasons for the omission of other pools of deadwood, litter and soils. The UNFCCC TA accepted the reasoning of RoS, considered the excluded pools are likely to be insignificant in the context of 2020 subnational FRL. However, it identified treatment of emissions particularly from soil, deadwood pools and fires as areas for technical improvement in future FRL submissions.

The NFI (2021) provides good quality data on above-ground and below-ground biomass carbon stock, disaggregated by stratum, land use and states (administrative units). These two pools are also considered the most significant in all national GHG inventories reported in national communications submitted by RoS and therefore, are included in this national FREL/FRL.

The estimation of the amount of dead wood in RoS is difficult, because significant amount of deadwood is collected directly by local communities living in the proximity of the forests to meet their immediate energy demands. Biomass energy is the main source for rural communities and for a significant portion of the urban population, it constitutes more than 60% of the energy balance in RoS. The Forest Product Demand Study (1995), estimated the amount of wood fuel directly collected at about 72% of total fuelwood used in RoS. This amount is mostly deadwood (fallen and stumps) and is not captured in available records and is consumed in the same year of its collection. Resources permitting, FNC needs to update the forest products demand survey of 1995, including household surveys, to develop quality estimates of the amount of wood directly collected from the forests.

RoS does not have repeated measurements NFI data on deadwood. The NFI (2021) provides the first data on the volume of dead wood remaining in the forests. In this submission RoS includes only standing deadwood in the estimation of emission associated with deforestation. According to the NFI field experts the data collected on standing deadwood is more reliable compared to the data on fallen and stumps deadwood.

The NFI (2021) does not provide information on soil, generally there is no good quality data on soil available in RoS, and this is also recognized in the GHGs inventory reported of RoS's national communications to date. Table 3 below summarize the carbon pools included in this submission.

**Table 3: The carbon pools included and omitted in the national FRL**

Pools	Inclusion in FRL	Justification
Aboveground biomass	Included	Significant, good quality NFI 2021 data
Belowground biomass	Included	Significant, good quality NFI 2021 data
Deadwood	Partially included	Standing deadwood, NFI 2021 data
litter	Not included	Not significant in drylands, Lack of data
Soil organic carbon	Not included	Lack of data

### **5.3. Greenhouse Gases:**

In the Sub-national Forest Reference Level (FRL) submitted in 2020, Sudan focused exclusively on carbon dioxide (CO<sub>2</sub>) emissions. The UNFCCC Technical Analysis (TA) identified the need to address non-carbon dioxide (non-CO<sub>2</sub>) gases (to maintain consistency with the national GHG inventory) as an area for future improvement. CO<sub>2</sub> remain the most significant gas in GHG inventory conducted in RoS (INC 2003, SNC 2013 and TNC yet to be published). The data and country specific parameters required to obtain good quality estimates of non-CO<sub>2</sub> are currently not available, RoS continue to use more default parameters, emission and conversion factors in most of the categories reported in its LULUCF inventory.

## 6. ACTIVITY DATA FOR DEFORESTATION

In the context of forest monitoring, Activity Data (AD) refers to information on human or natural actions that result in changes to forest cover and land use. Key activities tracked under AD include deforestation, afforestation, reforestation, and forest degradation. AD can be extracted from different sources, though for the current submission RoS relies preliminary on remote sensing information to track deforestation for an envisaged monitoring period of 2012-2021 at national scale. Whereas information on forest enhancement, including afforestation and reforestation, is mainly coming from national statistics records from Forest National Corporation (FNC). The current reporting is using an updated methodology for the second Forest Reference Level (FRL) submission considering the previous Technical Assessment (TA) team's review.

The stratified area estimate approach is an established method for assessing unbiased area estimates of land cover, land use and land use change, leveraging the statistical advantages of stratification to improve the precision around respective area estimates (Olofsson et al. 2014, FAO 2016).

For this FRL submission, two separate stratifications have been overlaid. The first stratification is based on ecological zones divide the forest landscape into distinct strata based on key variables such as forest type, land-use history, or ecological characteristics. This stratification has been employed in the NFI, and samples have been allocated to each stratum systematically, but with different intensities. It is further described in 6.1.1.

Further intensification on the above-mentioned stratified systematic grid has been added using a second stratification with low, medium and high probability of forest and forest change. This stratification is obtained through a forest change assessment based on satellite remote sensing. Subsequent sample allocation has been done by using a statistically optimized approach and samples have been placed in a random fashion. This second sampling strategy aimed at enhancing the precision specifically around the forest change area estimate and is further described in 6.1.2. The derivation of the stratification layer is outlined under 6.2.

The combined sampling design can be contextualized as a stratified random sampling and accordingly estimators have been employed as described in section 6.5.

## 6.1. Sampling Design

### 6.1.1 Systematic NFI grid based on permanent stratification

RoS implemented a systematic grid in 2017 as part of its National Forest Inventory (NFI) project to collect the field information at the national scale. Given the country's diverse ecosystems, a geospatial analysis was conducted to categorize RoS into distinct zones. This classification was based on available geospatial layers, including ecological zones, global tree cover, and land cover data. The resulting layer, referred as the strata boundaries, formed the foundation for generating the NFI grid. Table 4 provides a detailed description of the five strata.

**Table 4: Description of the Strata, along with total area covered by each stratum (Source: NFI Report Table 3.2)**


Stratum	Description	Area (ha)	Area (%)
I	The stratum that mostly comprised Deserts	67,674,160	36
II	The stratum characterized by semi-desert ecosystems (e.g. few Acacia trees and thorny bushes and xerophytes)	38,985,260	21
III	The stratum indicated as 'Low rainfall woodland Savannah' by Harrison and Jackson (1958)	35,972,310	19
IV	This stratum includes semi-arid, dry sub-humid, humid aridity zones. Forest and Woodland vegetation is mostly found here.	43,161,040	23
V	This stratum includes rivers and streams. It is probably the most heterogeneous since it is the stratum where human activities are dominated, and patches of vegetation (natural and not) found as riverine vegetation. This layer crosses all the latitudes of the country.	2,440,303	1

Recognizing the ecological diversity across the country, different sampling intensities were applied based on the specific characteristics of each zone defined in Table 5. For instance, the arid northern region, dominated by desert landscapes, was assigned a less dense grid with a higher expansion factor. In contrast, the forest-rich southern regions, characterized by dense vegetation, were assigned with a denser sampling grid (Figure 2). The distribution of sample intensities and field-collected samples for the NFI cycle (2017–2020) is summarized in Table 5.

**Table 5: NFI Sampling Design: Distribution of Sample by Stratum and total filed visited samples during NFI cycle 2017 -2020. (Source: NFI Report Table 3.2)**

Stratum No.	Total No. of sample units	Total sample collected in Field	Spacing of sample unit grid
Strata-I	107	-	80 * 80 Km <sup>2</sup>
Strata-II	123	20	80 * 40 Km <sup>2</sup>
Strata-III	431	159	40 * 20 Km <sup>2</sup>
Strata-IV	1,063	577	20 * 20 Km <sup>2</sup>
Strata-V	31	28	Random
<b>Total</b>	<b>1,755</b>	<b>784</b>	

The systematic grid is used as a basis for the AD collection as it covers the full country. It further facilitates consistent and repeatable data collection in the future, making it suitable for generating consistent long-term land use and land cover (LULC) statistics, as it is not subject to sampling variability. The activity data was collected across the entire grid population of 1,755 sampling units, though field data from NFI cycle is only available at 784 sampling units.



**Figure 2: Geographic distribution of sample intensities over each stratum**

### **6.1.2 Intensification of the sampling grid**

Stratum III and IV were subject to sampling intensification, as those are the ones containing forest. The intensification process was based on a second stratification that was derived through remote sensing based change analysis over the reference period using SEPAL, a cloud computing platform developed by FAO's forestry team. The approach follows a modified map subtraction workflow between two Forest/Non-Forest maps at the beginning and the end of the reference period. The important difference to a classical map subtraction is that it is not using the categorical output of the classifier, but rather the class probability of the pixels being forests. Subtracting those forest probabilities results in a more nuanced distinction that can be referred to as likelihood of being forest change. This layer of change likelihood is then used in a statistically optimized procedure to divide the area into relevant change strata and allocate an optimal number of samples randomly. The process to generate the stratification can be broken down into following steps:

1. Training data collection
2. Provision of satellite image composites
3. Creation of Forest non-forest (FNF) probability maps at time 1 and 2
4. Change probability layer (CP) through probability map subtraction
5. Unsupervised stratification of change probability layer
6. Optimal sample allocation

#### **6.1.2.1 Training data collection**

The training data was received from a prior land use and land cover mapping exercise conducted in 2020 by the Geospatial Department of the FAO. The training data was collected using eight Land Use Land Cover (LULC) classes namely: forest, bare rocks and soils, cropland, herbaceous, shifting cultivation, shrubland, urban areas, and water bodies. For this purpose, the data was remapped into two categories: forest and non-forest classes (Table 6). A total of 4,966 samples were collected in 2020 using Collect Earth Online (CEO) at a spatial scale of 100 meters using proportional allocation. Reclassification was conducted using a majority rule, where samples with greater than fifty percent forest are considered forest, and the remaining samples were assigned as non-forest. The training data was consistently applied across both years classification.

**Table 6: Categorization of training data into Forest and non-Forest class**

Sr. No.	LC class	Re-Categorization
1	Forest	Forest
2	Bare Rocks and Soil	
3	Cropland	
4	Herbaceous	
5	Shifting Cultivation	non-Forest
6	Shrubland	
7	Urban	
8	Water Bodies	

#### 6.2.2. Provision of satellite image composites

Wall-to-wall image mosaics generated from satellite data serve as a predictive input to the classification procedure alongside the labelled training data. In order to capitalize on the latest technological advances, RoS used a combination of data originating from optical and radar sensors. The selection was based on the availability of the data for the envisaged monitoring period and geographic coverage. For this analysis data from the Landsat missions, the Planet NICFI programme and the ALOS Kyoto & Carbon initiative was used.

**Landsat Composite:** Landsat composites for the year 2011 and 2023 were created using SEPAL's Optical mosaic recipe. First, an annual composite for the year 2011 was generated using satellite data from Landsat 4-5. Missing areas of incomplete coverage were filled in by data from the preceding years. Similarly, Time-2 composites were generated for the year 2023 and 2024 using step by step guide available at SEPAL documentation [page](#). The year 2024 is introduced as a buffer year, the idea behind temporal buffer is to avoid potential omission of changes that might happen at the start or end of year. Such changes would not be captured in the subsequent change analysis when using the best pixel composite, since pixel information in these mosaics could come from any point in time within selected dates. The buffer year is defined here as start of envisaged monitoring period and minus one year (2010 & 2011) and end of monitoring period and plus one year (2023 & 2024). For 2011 it was not possible to create two separate mosaics for year 2011 and 2010 because of limited data availability therefore, a combined composite for the years 2009-2011 was created.

**Planet Composite:** The annual composites are generated by aggregating data from monthly basemaps of Planet NICFI. The mosaics are extracted using Planet mosaic recipe in SEPAL for the year 2023 and 2024. A step-by-step process is available at the [link](#).

**ALOS Mosaic:** L-Band Radar data for the envisaged monitoring period was available through ALOS Kyoto and Carbon (ALOS K&C) initiative, which provides global 25m PALSAR/PALSAR-2 mosaics on an annual basis. Backscatter data from long wavelength radar systems provides complementary information on the land cover. As the backscatter over vegetated areas is coming from woody elements, it is less dependent on the presence of leaves and can therefore detect seasonal dry forests outside their leave-on period.


The ALOS K & C mosaic were created using the following Google Earth engine (GEE) script. This code creates a mosaic over RoS for the year 2010 and 2023 separately. The workflow defines the area of interest (boundary), retrieved the composite for specified year, and applies noise reduction and calibration. The key metrics like texture (contrast, entropy) and polarization ratios (HH/HV) are computed to analyse surface properties. The processed image is visualized and exported as an asset to GEE. The script is very automatic and only required two changes namely AOI and year.

**Data Retrieved:** All mosaics were exported to GEE as an asset at 30-meter scale regardless of original resolution. This was done to maintain the consistency among datasets used for activity data (AD). It's important to mention that stratification layer leverage the data from buffer year, but reference data collection was strictly confined to 2011-2023.

### 6.2.3. Forest Non-Forest (FNF) Probability Maps

Forest/Non-Forest probability maps were produced using supervised classification in SEPAL, where the annual composites from the relevant years were combined with the training data to generate FNF probability maps for each of the respective years. The classification was run separately for both 2011 and 2023 using the stable set of training samples. For the FNF probability map 2011 input data from ALOS K & C and Landsat was used whereas for 2023 ALOS K & C , Landsat and Planet composite from respective and buffer year were used for classification. Table 7 is providing detailed description of input images along with band combinations and indices used for classification, All datasets are downloaded as GEE assets and asset id are provided in Annex-5. A random forest classifier was used which was calibrated using input training data and annual composites from respective years. One of the important parameters using random forest classifier is the number of trees used in the underlying bootstrapping. By default, this is set to 25 and was increased to 1000. Instead of the categorical output, the class probability layer of forest is extracted

with values ranging from 0-100. The resulting classification was exported to GEE. The details workflow of this analysis is given in Figure 3 of the document.



**Figure 3: Showing workflow of the activity data**

#### 6.2.3.1 Post-Processing

The coverage of the ALOS mosaics was incomplete due to small gaps. To fill the gaps, an additional forest probability layer was generated using only the optical data, following the same process as explained above. Gaps in the multi-sensor probability layers were then filled in with the probability values of the classification using only optical data.

**Table 7: Depicting details of Optical and Radar composite used for classification along with relevant bands.**

Sensor	Monitoring Time	Composite Image Sensor	Band & Index used	Composite Year
Optical	Time- 1	Landsat 4-5 composite	Red, NIR, SWIR-1, SWIR-2, SAVI, NDFI	2009 -2010 <sup>1</sup>
Radar		ALOS K & C Timescan	HH, HV, HHV_ratio	2010
Optical	Time -2	Landsat 8	Red, NIR, SWIR-1, SWIR-2, SAVI, NDFI	2023
		Landsat 8		2024

<sup>1</sup> Because of insufficient image availability composite image period was extended to 2009-2010

		Planet NICFI Basemap	Blue, Green, Red, NIR	2023
		Planet NICFI Basemap		2024
Radar		ALOS K & C Timescan	HH, HV, HHHV_ratio	2023

#### 6.2.4. Change probability Layer

After generating the FNF probability layers for the start and end of the envisaged monitoring period, the change probability layer was generated by subtracting the FNF probability of 2011 with the one from 2023. As an increase in forest probability would lead to negative values, the operation is using the absolute values of the difference, thus leading to the change probability layer with values ranging from 0 to 100 as follows:

$$\text{Change Probability layer} = \text{Abs} (\text{FNF probability 2011} - \text{FNF probability Map 2023}) \quad \text{Equation- 1}$$

##### 6.2.4.1. Inclusive Forest Mask

The northern part of the RoS is largely desert, with no indication of forest, as confirmed by the initial interpretation using systematic grid sampling. Therefore, prior to stratification, a potential non-forest area was masked out by applying a threshold value of five. This threshold was determined through visual interpretation of Forest and Non-Forest (FNF) probability maps for both 2011 and 2023. The rationale for setting this threshold was to generate an inclusive forest mask that captures the potential forest extent, ensuring that no forest areas or subsequent changes were overlooked.

#### 6.2.5. Stratification of change probability

KMeans is a clustering algorithm that groups data points based on statistical assumptions, whereby it minimizes the intra-cluster variance while maximizing inter-cluster differences. Therefore, this method is effective in stratifying skewed populations (Kozak (2011)). It has been used on top of the masked change probability from the previous step, grouping the data into 3 strata.

#### 6.2.6. Sample Allocation

Sample allocation was based on the optimal sample allocation scheme following Neyman (Neyman, 1934, Cochran, 1977). This method uses the stratum size and the underlying variability

in change to calculate optimal number of samples to be allocated in each stratum. Strata with larger populations or higher variability get more samples. It is calculated as follows:

$$n_h = \frac{N_h \cdot S_h}{\Sigma(N_h \cdot S_h)} \cdot N \quad \text{Equation -2}$$

$n_h$  = Number of samples allocated to stratum  $h$

$N_h$  = The size (or area) of stratum  $h$  (e.g., total count or area in hectares)

$S_h$  = The standard deviation of the variable of interest (e.g., change probability) within stratum  $h$

$\Sigma(N_h \cdot S_h)$  = The weighted variability across all strata (used to normalized)

$N$  = Total number of samples to be collected (target sample size)

In case of RoS, three strata were created within a mask generated after applying a threshold whereas area outside mask was considered as fourth stratum. The sample intensification using stratification was performed only in permanent strata (NFI Stratum) three and four of the country. This decision was taken based on preliminary analysis of systematic grid which confirmed the proposition of no forest loss in strata one and two whereas strata five was skipped because most of the activities around Nile River (Strata-5) are cyclic in nature and are part of local forest management practices. Table 8 is showing sub-stratification of strata three and four.

**Table 8: Sample intensification by stratum along with total area of stratum in hectares**

No.	NFI Strata	Temporary Strata	Allocated Sample	Total area
1	Stratum I	Masked	107	67,674,160
2	Stratum II	Masked	123	38,985,260
3	Stratum III	Outside mask / no change	1,055	27,467,985
		Low change probability	551	6,735,925
		Medium change probability	213	1,516,955
		High Change probability	83	251,445
4	Stratum IV	Outside mask/no change	183	2,109,760
		Low change probability	2,130	23,631,535
		Medium change probability	1,940	13,277,357
		High Change probability	1,142	4,142,388
5	Stratum V	Masked	31	2,440,303
<b>Total Sample</b>			<b>7,558</b>	<b>188,233,073</b>

### 6.3. Data Collection

A customised response design was prepared for data collection (Figure 4), which was later translated into a Collect Earth (CE) survey for data collection. Collect Earth is an open-source tool

developed by the FAO under the Open Foris initiative, designed to simplify land use, forestry, and environmental assessments through satellite imagery and geospatial data. It is integrated with platforms like Google Earth and Google Earth Engine, which allow users to monitor deforestation, afforestation, and land-use changes with ease. It has been widely used for reference and monitoring data collection by various countries. The response design used for data collection is attached as an Annex-8 of this document which provides definition of each LULC and changes classes along with explanatory variables.


Data was collected for the period 2011 until 2023 in a backward moving direction. First the information on Land Use Land Cover (LULC) class was collected (Figure-5, survey card one) for 2023. In case of forestland, an additional question about tree cover percentage needed to be answered. In case of a change event, information about LULC changes was collected in a second survey card. This information includes the respective year and the driver of change. A third survey card gathered general information over each sample such as satellite imagery used for decision making, and confidence on collected information along with comments.

## Reference Data collection Survey Design for Sudan, FRL 2024


### 1- LULC 2023 (one class to be selected, in case of Forest Land questions could be will be asked, as below)



### 2- LULC Changes (Based on class selected, further questions are asked, as mentioned in text box in numeric order)



**Figure 4: Customized Response Design established for Republic of Sudan for data collection**



**Figure 5: Collect Earth Survey Card used for activity data collection using Collect Earth software**

#### 6.4. Capacity Building

Two capacity building workshops were organized by FAO with RoS Team to develop the capacity of the team. In the first training workshop expert from FAO familiarizes the participants with collect earth software and response design to be used for data collection. After the workshop samples were equally distributed among the participants. The workshop was followed by multiple troubleshooting sessions with the team during the data collection process. After the first round of data collection an additional set of samples were generated using the stratification layer and distributed among team members, list of core Remote Sensing team is provided in Annex-9 of this document. The second training workshop conducted by FAO was designed to familiarize the team with the SEPAL platform, focused on two primary processing recipes: 1) Optical Mosaic and 2) Planet Mosaic. Following the workshop, the team was assigned the task to generate Landsat and Planet composite images for envisaged monitoring period of 2011 -2023, which were then used to produce FNF probability maps.

#### 6.5. Results

After data collection area estimates are calculated at Strata and States levels, Strata served at permanent stratification for national forest monitoring system. Whereas forest management is mainly governed by state administration. Both level estimates are extracted from same set of

samples, where information of strata and states is extracted by spatially overlaying samples over the respective shapefiles.

$$A_{ij} = \frac{n_{ij}}{n_i} \cdot A_i \quad \text{Equation- 3}$$

Where:

$A_{ij}$  = Area of class  $j$  in stratum  $i$ .

$n_{ij}$  = Total number of samples of class  $j$  in stratum  $i$ .

$n_i$  = Total number of samples in stratum  $i$ .

$A_i$  = Total area of stratum  $i$ .

$$SE = \sqrt{\frac{p_{ij} \cdot 1 - p_{ij}}{n_i - 1}} \cdot A_i \quad \text{Equation - 4}$$

Where:


$SE$  = Standard Error calculated for each class  $j$  in stratum  $i$

$p_{ij}$  = Proportion of class  $j$  in stratum  $i$ , calculated as  $p_{ij} = \frac{n_{ij}}{n_i}$

$n_i$  = Total number of samples in stratum  $i$ .

$A_i$  = Total area of stratum  $i$ .

The areas of each class within a stratum is calculated using equation 3, where the number of samples in each class within the relevant stratum is divided by the total number of samples in that stratum and then multiplied by the total area of the stratum. The Standard Error (SE) for a given class in a stratum is calculated using the equation 4, which is then multiplied by the Z-score (1.64) to derive the 90% confidence interval (CI) and converted to percentage. The data collected pertains to six classes: forest loss, forest degradation, forest gain, other changes, stable forest and stable non forest. Detailed description of each class is provided in the data collection sections and Annex-10 of this document. RoS reports forest loss from this analysis for the current Forest Reference Level (FRL) submission. During the reference period (2012 – 2021) the total area of deforestation observed is estimated at 2,059,749 ha, with an annual deforestation rate of 205,979 ha over this period.



**Figure 6: Annual deforestation trend from 2011-2023**

Figure 6 illustrates annual deforestation rates over the reference period, does not clearly indicate a specific trend. Though, the highest deforestation rates were recorded in year 2014, followed by 2013 and 2019, while the lowest deforestation occurred in 2022. Operators often struggled to pinpoint the exact year of deforestation due to inconsistent availability of high-resolution data, particularly during the early years of the monitoring period. Consequently, the accuracy of determining the precise year of change heavily depends on the availability of high-resolution data.

### 6.5.2. Results by States


RoS is divided into eighteen states, with forests present only in thirteen of them, predominantly located in Strata 3, 4, and 5. According to the current analysis, the forest changes are mainly confined in strata 3 and 4 of the country, which are main forest vegetation zones. The stratum four, forest rich zone in the southern part of the country, exhibits the highest levels of deforestation, driven predominantly by agricultural expansion and other human activities. Whereas Stratum three, also shows significant changes, though at lower rate compared to stratum four. Cyclic forest loss identified mainly in strata five along the Nile River and its tributaries and this is primarily part of sustainable forest management practices in RoS. Strata one and two are the desert and semi-desert area, only one sample was identified as a forest. Another significant activity observed is forest degradation the main driver of degradation is wood fuel and overgrazing activities. Table 9 provides area estimates derived from the current analysis at the state level. The analysis shows that Southern Kordofan exhibits the highest rates of deforestation and forest gain, followed by Southern Darfur. In contrast, the lowest rates of forest change are recorded in Western Darfur. no forest presence or related changes were observed in the Northern States, Red Sea, Gezira, Khartoum and River Nile states. The most significant forest degradation was observed in Western Kordofan,

followed by Southern Kordofan and Darfur. The overall confidence intervals for state-level estimates vary widely, ranging from 11% to 71%. These estimates are calculated using the equations outlined in section 5.4.

**Table 9: Stratified Area Estimates by States in Hectares**

Name	Forest Degradation	Forest gain	Forest loss	Other changes	Stable forest	Stable non forest	Total area (ha)
<b>Blue Nile</b>	58,477	75,565	102,318	21,977	1,167,525	2,382,186	3,808,048
<b>Cantral Darfur</b>	29,033	82,226	52,980	26,251	900,231	1,901,420	2,992,140
<b>Eastern Darfur</b>	94,222	166,865	158,116	123,075	1,935,964	3,801,227	6,279,468
<b>El Gadarif</b>	21,566	17,593	76,087	17,726	188,717	5,551,701	5,873,390
<b>Gezira State</b>	-	-	-	-	-	2,355,693	2,355,693
<b>Kassala</b>	3,029	11,095	74,914	6,657	13,688	5,242,495	5,351,878
<b>Khartoum State</b>	-	-	-	-	-	2,137,267	2,137,267
<b>Northern Darfur</b>	3,627	48,435	33,035	98,620	1,134,391	31,152,713	32,470,821
<b>Northern Kordofan</b>	-	10,151	185,558	38,261	202,037	17,108,146	17,544,153
<b>Northern State</b>	-	-	-	-	-	34,710,102	34,710,102
<b>Red Sea</b>	-	-	-	-	-	22,786,131	22,786,131
<b>River Nile</b>	-	-	-	-	-	13,210,477	13,210,477
<b>Sinnar</b>	43,330	60,645	52,357	11,095	575,684	2,997,066	3,740,176
<b>Southern Darfur</b>	151,610	188,496	424,989	98,392	3,068,465	4,322,995	8,254,948
<b>Southern Kordofan</b>	184,741	246,518	745,063	109,176	2,417,266	4,373,739	8,076,503
<b>Western Darfur</b>	17,751	23,320	33,791	26,036	375,170	1,948,085	2,424,153
<b>Western Kordofan</b>	206,535	147,639	423,512	300,529	2,594,935	8,053,002	11,726,152
<b>White Nile</b>	6,657	18,004	22,402	50,486	81,360	4,312,664	4,491,572
<b>Grand Total</b>	820,579	1,096,551	2,385,119	928,281	14,655,434	168,347,109	188,233,073

The majority of forest-related changes are concentrated in Southern and Western Kordofan followed by Southern Darfur and Eastern Darfur.



**Figure 7: Showing trend of forest changes by states over 2012 -2021**

Tables 10 and 11 below provide detailed annual estimates of deforestation (forest loss) and forest degradation by state.

**Table 10: Annual Areas of Deforestation by State in Hectares**

Year	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	Total
Blue Nile	-	-	7,255	21,353	10,471	14,099	3,627	10,471	7,255	20,943	6,844	-	-	102,318
Central Darfur	-	21,566	6,844	13,688	-	-	-	3,627	3,627	-	3,627	-	-	52,980
Eastern Darfur	20,532	21,566	17,939	11,095	32,880	3,627	3,627	6,657	-	18,349	14,722	-	7,122	158,116
El Gadarif	3,627	-	6,844	-	7,255	-	10,471	3,627	26,947	13,688	-	-	3,627	76,087
Kassala	12,225	-	27,557	3,627	7,122	6,657	-	-	-	10,471	3,627	3,627	-	74,914
Northern Darfur	-	-	-	12,225	6,844	13,966	-	-	-	-	-	-	-	33,035
Northern Kordofan	6,844	10,882	3,627	12,225	-	6,844	-	29,663	7,255	22,376	-	-	7,122	106,838
Sinnar	6,844	-	-	20,943	-	-	-	3,627	78,719	3,627	6,844	-	10,471	131,076
Southern Darfur	66,789	18,783	64,287	94,559	14,099	20,943	36,288	10,749	20,943	20,943	24,570	10,471	21,566	424,989
Southern Kordofan	36,075	53,588	114,795	65,717	50,584	56,607	86,051	75,154	77,550	53,588	32,235	18,137	24,981	745,063
Western Darfur	12,225	-	-	-	-	10,471	-	-	-	-	11,095	-	-	33,791

Western Kordofan	18,783	64,507	26,734	96,310	18,349	47,266	10,471	3,627	23,972	64,763	20,532	3,627	24,570	423,512
White Nile	3,029	-	-	3,029	-	-	-	3,627	3,627	3,029	3,029	3,029	-	22,402
Grand Total	186,974	190,892	275,882	354,771	147,604	180,480	150,536	150,831	249,895	231,778	127,126	38,892	99,459	2,385,119

**Table 11: Annual Areas of Forest Degradation by State in Hectares**

Year	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	Total
Blue Nile	-	11,095	-	11,095	6,844	-	-	3,627	11,095	11,095	-	-	3,627	58,477
Central Darfour	6,844	11,095	-	-	-	-	-	-	-	11,095	-	-	-	29,033
Eastern Darfour	22,189	-	-	-	-	17,939	-	-	7,122	40,128	6,844	-	-	94,222
Gadarif	-	-	-	-	-	-	-	-	6,844	3,627	11,095	-	-	21,566
Kassala	-	-	-	-	-	-	-	-	-	-	3,029	-	-	3,029
Northern Darfour	-	-	-	-	-	-	-	-	-	3,627	-	-	-	3,627
Sinnar	-	3,627	-	3,627	-	6,844	-	-	14,722	7,255	3,627	-	3,627	43,330
Southern Darfour	-	11,095	35,254	36,911	-	6,657	-	11,095	14,722	24,783	11,095	-	-	151,610
Southern Kordofan	-	35,254	7,255	42,919	6,844	10,882	27,787	14,099	10,471	3,627	14,722	3,627	7,255	184,741
Western Darfour	-	-	-	-	-	3,627	-	11,095	-	-	-	-	3,029	17,751
Western Kordofan	21,566	6,844	24,783	64,287	-	6,844	39,915	6,844	17,315	7,255	3,627	-	7,255	206,535
White Nile	-	3,627	-	-	-	3,029	-	-	-	-	-	-	-	6,657
Grand Total	50,599	82,636	67,291	158,839	13,688	55,822	67,702	46,759	82,291	112,491	54,039	3,627	24,793	820,579

Table 12 presents a breakdown of deforestation classes by land use and land cover (LULC) categories. The analysis indicates that agricultural expansion is the primary driver of deforestation in Sudan, followed by a "other" class. The current assessment focused on collecting data related to the following key drivers: agricultural expansion, infrastructure development, petroleum exploration, mining, and an additional category labelled as "other." This latter category was used when interpreters were unable to classify a driver within the predefined list.

The list of drivers was developed based on findings from the national study on deforestation and forest degradation drivers (DoDD 2018). As a result, further disaggregation of the "other" category is not possible due to the limited specificity in the data. However, a review of comments associated

with the "other" category suggests that many of these samples are linked to grazing on agriculture lands. In unspecified cases, the interpreters did not provide sufficient detail to enable a more precise classification.

**Table 12: Disaggregation of deforestation area by LULC classes**

Forest Drivers	Forest Loss (2011-2023)	LULC
Agriculture Expansion	1,772,616	Cropland
Infrastructure Development	10,471	Settlement
Others (Heavy Grazing & others)	594,777	Grassland + other land
Petroleum Exploration	7,255	Settlement

## 6.6. Quality Management

For quality management, five percent of samples from systematic grid were selected randomly by stratum for multi-interpretation. These samples were distributed among all participants, list of participants is provided in the Annex 9 of this report. After the data collection, analysis was performed in python where a customized script was developed for analysis. That mainly focused on agreement analysis among operators by LULC classes and change classes. Results were compared with overall average of aforementioned classes. The analysis showed a great variability among participants for both LULC and forest change classes. The performance was analysed by identifying top five interpreter with majority agreement for both LULC and forest change classes. The analysis showed that the overall average of LULC classes were found very close to the top five interpreters (Figure 9). However, the average of the top five interpreters in forest change classes was very low compared to overall average. The one conclusion that can be drive from this analysis is that change classes are generally very hard to interpret compared to LULC classes.

**1- LULC Classes 2023 - Average**

Class	Average
Cropland	22.50
Forest Land	16.50
Grassland	26.38
Other land	17.81
Plantation	2.50
Settlements	1.14
Wetland	1.09
(blank)	6.00
Grand Total	

**2- First five Interpreters with majority agreement by LULC 2023**

	Operator - 1	Operator - 2	Operator - 3	Operator - 4	Operator - 5	Average
Cropland	24	24	23	18	24	22.60
Forest Land	17	17	17	17	21	17.80
Grassland	29	29	30	28	26	28.40
Other land	13	13	13	20	15	14.80
Plantation	2	2	2	1		1.75
Settlements	1	1	1	1	1	1.00
Wetland	1	1	1	2		1.25
Grand Total	87	87	87	87	87	

**3- Forest Change Classes 2023 - Average**

Class	Average
Forest Degradation (F - F)	2.42
Forest Enhancement (F-F / NF - F)	3.38
Forest gain	2.50
Forest Loss (F - NF)	1.13
No Changes	73.13
Other changes	7.69
(blank)	3.50
Grand Total	

**4- First five Interpreters with majority agreement by Forest Change Classes**

Class	Operator - 1	Operator - 2	Operator - 3	Operator - 4	Operator - 5	Average
Forest Degradation (F - F)		1	1	1	1	1.00
Forest Enhancement (F-F / NF - F)						
Forest gain	1		1	1	1	1.00
Forest Loss (F - NF)	1	1		1		1.00
No Changes	81	81	80	78	79	79.80
Other changes	4	4	5	6	6	5.00
Grand Total	87	87	87	87	87	

**Figure 8:** Showing comparison of top five interpreter average with overall average by LULC and forest change classes. Table 1 refers to LULC overall average by classes, Table 2- is showing average of top five interpreter by LULC classes. Table 3- is showing forest change classes overall average. Table 4 – showing an average of first five interpreters by forest change classes.

**Agreement Matrix:** Table 13 presents a matrix illustrating the agreement among interpreters by LULC classes. The fractions in each column represent the agreement among interpreters, with "MOD" indicating the majority class identified during interpretation. The highest agreement is observed for the cropland class, where 82% of interpreters agreed. Whereas the agreement for forestland is 72%, as interpreters often found it challenging to distinguish between plantations and natural forestland. This difficulty arises because plantations in RoS are typically random in nature and lack the structured patterns observed in other countries, making them harder to differentiate from natural growth.

**Table 13: Matrix of by LULC classes with majority agreement**

Value Mode	Cropland	Forest Land	Grassland	Other land	Plantation	Settlements	Wetland
<b>Cropland</b>	0.817708	0.010417	0.104167	0.044271	0.020833		0.002604
<b>Forest Land</b>	0.031915	0.719858	0.230496		0.01773		
<b>Grassland</b>	0.040302	0.130982	0.715365	0.108312		0.005038	
<b>Other land</b>	0.066176		0.113971	0.8125	0.003676	0.003676	
<b>Plantation</b>	0.125	0.25			0.625		
<b>Settlements</b>	0.0625			0.0625	0.0625	0.8125	
<b>Wetland</b>		0.0625	0.0625	0.1875			0.6875

## 7. Description of the National Forest Inventory (NFI):

RoS's NFI (2021) is the main source of emission factors for deforestation. FAO through the technical assistance agreement with the World Bank and FNC has provided the technical support capacity resources for the NFI, NFMS/MRV and other related activities of the REDD readiness programme. The design used for the NFI is one that promoted by FAO and used in several countries. The field measurements of NFI (2021)<sup>2</sup> were conducted during the period 2017-2019 and the report was published in 2021.

### 7.1. NFI Sampling Design:

A systematic cluster sampling design was adopted for the NFI in RoS. Sampling Units were selected at the grid intersection of the latitude longitude degrees. The number of SU to be surveyed was determined by the available financial and human resources. Stratification was adopted as ecological zones were assumed to improve the design. RoS area was stratified into five strata. Sample Units (SUs) were allocated to each stratum according to vegetation density as shown in Figure 2 above. Initially, a total of 1755 SUs were planned, then reduced to 968, and finally 784 SU were visited, the rest were inaccessible due to security or physical reason.

Data is collected in the field through observations, measurements and interviews at different levels within the limits of the sampling units (SU) and in smaller subunits, the plots, subplots, Land Use/Cover Sections (LUCS) and Land Use/Cover Classes (LUCC) demarcated within the sampling units, see Annex 3.

- A **sampling unit (SU)** or “Tract” is a square surface area of 1 km x 1 km. The coordinates of the south-west corner of the SUs correspond to those of the points selected in the systematic sampling frame. Each SU contains **four** field plots, which means in total 3136 (784x4) plots have been surveyed in this NFI cycle.
- The **plots** are rectangles, with surface areas measuring 20 m wide and 250 m long (area 0.5 ha) within the SU. They start at each corner of an inner 500 m square (same centre as SU's), and are numbered clockwise from 1 to 4.

Each plot is divided into Land Use/Cover Sections (LUCS) identified in the field as shown in the example below. Data related to grazing, cropping and forest characteristics, management and


---

<sup>2</sup> Full description of NFI (2021), is found at:

[https://www.fao.org/fileadmin/user\\_upload/faoweb/Themes\\_pages/Forests/REDD-NFM/RoS\\_MRV/RoS - NFI\\_Report.pdf](https://www.fao.org/fileadmin/user_upload/faoweb/Themes_pages/Forests/REDD-NFM/RoS_MRV/RoS - NFI_Report.pdf)

resources use and users are collected within the LUCS.

**Figure 4:** An example of Land Use/Cover Sections (LUCS) distribution within a plot



**Figure 9: Land use Land Cover Section**

## 7.2 Data Collection Process

Data are collected by the field teams for SUs, plots, subplots, measurement points, land use/cover section (LUCS), land use/cover class (LUCC) and interviewees. The main information sources for the assessment are:

- Field measurements and observations.
- Interviews with key informants (external and internal), focus groups and individuals and randomly selected households.

Different variables are collected depending on data collection levels:

- **Plot:** identification of different land use/cover sections (LUCS) and measurements of trees and stumps with a  $\text{Dbh} \geq 20 \text{ cm}$  in forest, or  $\geq 10 \text{ cm}$  for the trees outside forest. A plan of the plot indicating in particular land use/cover sections limits must also be completed.
- **Land Use/Cover Section (LUCS):** corresponds to the land use/cover sections identified along the plot. Information collected at this level includes: general information related to the LUCS; forest and other wooded lands management practices (harvesting, silviculture, etc.); and structure, as well as crop management practices.
- **Land Use/Cover Classes (LUCC):** corresponds to each land use class found in the SU (in all 4 plots). Information on forest and trees, on environmental services, pests and diseases, invasive and threatened species, wildlife abundance, and land use change is collected at this level.
- **Rectangular Subplot (RSP):** shrubs (in all LUCC), trees with  $20 \text{ cm} > \text{Dbh} \geq 10 \text{ cm}$  trees (only in forest LUCC), indicator plant species and non wood forest products (NWFP) are inventoried at this level.
- **Circular Subplot (CSP):** data on tree regeneration ( $\text{Dbh} < 10 \text{ cm}$  and height  $\geq 1.30 \text{ m}$ ) data are collected at this level (only in forest, OWL and woodlots) .
- **Litter Subplot (LSP):** at this level, data is collected on litter, which includes all non-living biomass with diameter less than 10 cm.

- **Fallen Deadwood Transect (FDT):** measurements of fallen deadwood branches ( $\geq 10$  cm) are done along the transect lines.
- **Measurement point (MP):** topographic and soil data is collected at the three measurement points.

### 7.3. Data Analysis

The information of the NFI field survey was stored in files (e.g. trees  $>10$  cm, small trees  $<10$  cm, stumps, etc) using the Open Foris Collect platform. Silva Metricus software has been used to perform calculations based on these files, Annex 4 provide the description of the NFI data analysis, calculation and the procedures used in driving the estimates of the NFI data including the volumes estimates used in this submission.

During the fieldwork stage, every effort was made to minimize measurement errors through training processes for the crew members, as well as the use of the best available measurement instruments. This is in addition to the review and quality checks performed twice for data entry, cleansing, transition to software and recalculations. The software Silva Metricus was used for the calculations of the sampling errors following the approach of “ratio estimator”.

At the national level, the overall variability of the NFI is very low, e.g. 9.11% for the forest land, because all the 784 Sampling Units are considered. However, when the calculation is disaggregated to State level, the number of sampling units included in the calculation of each State decrease and therefore the variability increase (see Table 15 below).

## **8. Emission Factors for Deforestation:**

### **8.1. Revised estimates of above-ground biomass:**

As indicated in section 7.4 and 14 of the initial FRL submission (January 2025), the emission factors for estimating emissions from deforestation were desired to be obtained from the published data of the NFI (2021). However, as explained in the NFI report, during field surveys teams could not visit all planned sample units (SUs). Initially, a systematic grid with different intensities was designed with 1755 SUs. A visual assessment of these SUs was performed using Collect Earth removing SUs without vegetation cover, resulting in a total of 968 SUs planned for field measurements. However, only 784 SUs (81%) were visited and measurements of trees collected from them since the rest of SUs were inaccessible due to security or physical reasons. This inaccessibility results in a violation the proposed sampling design of the NFI, thereby challenging strata-wise, state-level or any area- wide assessments in the country.

As part of the FRL project, there was a plan to improve the NFI data and hence the EFs for estimating the FRL. The original plan was to undertake field measurements from the unvisited SUs. However, because of the current war situation this was also not possible. RoS Sudan requested support from the University of Maryland through FAO to explore using the Global Ecosystem Dynamics Investigation (GEDI), to fill in gaps in the NFI data due to the un-visited sample units. An initial report of this collaboration containing the description of the methodology used and initial estimates of aboveground biomass density (AGBD) by state is included in annex 7 of the FRL submission. The methodology uses two sources of Earth Observation (EO) data – i.e. the GEDI data and the Forest Probability Map as auxiliary data to support the NFI. Note, the EO datasets are calibrated to the visited NFI plots, in a geostatistical model. The EO datasets serve as layers that augment the NFI estimates of AGBD. The results thereby allow predictions of forest AGBD with negligible systematic error. Based on the EO-supported model-based predictions of AGBD revised EFs have been developed. As compared to the Annex 7 of initial FRL report, changes are made in the manner in which model predictions, per state, were obtained:

1. Model-based predictions of AGBD are made for all locations in the original planned NFI grid, but the forest biomass estimates only use predictions at locations which are visually interpreted as stable forest, degraded forest or forest gain.
2. For some SUs, there is a possibility that the Forest Probability map or the GEDI data is no-data (NA). Further, some poor quality EO data pixels (due to clouds, for example) can lead to very high and unrealistic model-based predictions of AGBD. In case AGBD values larger than 500 Mg/ha were predicted at any location, they were removed from the prediction samples.

The methodology and results are described in the sections that follow.

### 8.1.2. Methodology

Sudan's National Forest Inventory (NFI) provides estimates needed for vegetation and forest assessments of the country, including volume and aboveground biomass density (AGBD). The sampling design of the Sudanese NFI is a systematic grid, with its density varying by ecological strata across the extent of the country. Spatial gaps in the NFI, however, are present due to the inaccessibility of some areas of the country. These spatial gaps from unvisited plots result in a violation the proposed sampling design of the NFI, thereby challenging strata-wise, state-level or any area-wide assessments in the country. Visited and measured NFI plots are displayed in Figure 1a.




Figure 10: NFI estimates of aboveground biomass and volume, and the covariates used in the geo- statistical model. The covariates are NASA GEDI-estimated aboveground biomass and the Forest Non-Forest (FNF) Probability Map 2023 over Sudan.

To overcome this challenge, Earth Observation (EO) datasets are used to provide supplementary information in support of the ground-based inventory. In a model-based

framework, for example, the use of auxiliary information from EO datasets can allow spatial scaling of the NFI estimates, thereby filling gaps where NFI plots were not visited, allowing area-wide estimations of aboveground biomass and improving their precision accuracy (e.g., [Babcock et al., 2018](#); [Breidenbach et al., 2021](#); [Emick et al., 2023](#); [May et al., 2023](#)). Important to note is that, in the case of Sudan, EO datasets are not replacing or recalibrating (or thereby biasing in any way) the NFI-estimated vegetation biomass. The EO datasets simply serve as layers that augment, enhance, support and supplement the NFI estimates of vegetation biomass. Irrespective of the pixel-level accuracies of the supportive EO map layers, they can be modeled to augment the NFI as long as they capture spatial variations across the domain of the country. The other important considerations for the use of auxiliary EO datasets to enhance the NFI are (1) the visited NFI plots provide a fair representation of the range of ecoregions and vegetation AGBD found over the extent of Sudan, and (2) the auxiliary EO datasets are acquired or created independent of the NFI data.

For the case of Sudan, estimates of vegetation aboveground biomass from the NASA Global Ecosystem Dynamics Investigation (NASA GEDI) ([Dubayah et al., 2020](#)) and the Forest Non-Forest (FNF) Probability Map 2023 are used as auxiliary EO data sources. The former dataset, i.e. the NASA GEDI data, is available as 25-m footprints of aboveground biomass estimated from high resolution laser-measured vegetation height, collected between years 2019-2023 ([Dubayah et al., 2021](#)). These footprint-level AGBD estimates are based on generic global models ([Duncanson et al., 2022](#)), and their accuracy over Sudan is hitherto unknown. However, note again, we are interested in only using this GEDI data to augment the NFI, hence calibrating it to the NFI-estimated AGBD, irrespective of its footprint-level accuracy. The GEDI footprint-level AGBD estimates are averaged in a 1 km × 1 km grid across Sudan, making a near-continuous map of estimates (Figure 1b). The latter dataset, i.e., Forest Non-Forest (FNF) Probability Map 2023 (Figure 1c), is described previously in sections of Activity Data for Sudan.

#### **8.1.2.1. Geostatistical model relating EO and NFI data**

The method adopted for Sudan is based on the geostatistical approach provided in [Hunka et al. \(2025\)](#). This section provides a brief overview of the approach, as well as the link to access to the open-source R code released to run the model for the case of Sudan.

A geostatistical model-based approach is used to relate the Sudanese NFI to the auxiliary EO datasets, with inference conducted through Bayesian methods. A geo- statistical model is

effective in capturing spatially continuous variables (here, AGBD) that have been measured only at a finite number of sites (here, the visited NFI locations). The model itself is a linear regression, relating the NFI-estimated aboveground biomass to the EO datasets' pixel values at the locations of the NFI plots. EO dataset values are extracted as the mean value of EO map pixels that intersect an NFI plot cluster (of size 500 m × 500 m), weighted by the fraction of the pixel that is covered by the cluster (see ‘[exactextractr](#)’ package in ‘[R](#)’). In the linear regression, however, we further add a geostatistical component, such that components of the model parameters are allowed to vary across the domain of Sudan. Adding such components allows to account for spatial variation in the regression coefficients and spatial autocorrelation in the model residuals ([May et al., 2023](#); [Emick et al., 2023](#); [Hunka et al., 2025](#)). We assume the model

$$y(s) = (\alpha + \tilde{\alpha}(s)) + (\beta + \tilde{\beta}(s))x_1(s) + (\eta + \tilde{\eta}(s))x_2(s) + \epsilon(s), \quad (1)$$

where

$s$  is a two-dimensional coordinate vector defining a spatial location

$y(s)$  is the NFI-estimated AGBD at location  $s$

$x_1(s)$  is the GEDI estimate of vegetation biomass at location  $s$

$x_2(s)$  is the forest probability estimate at location  $s$

$\alpha$  is the intercept

$\beta$  is a model parameter

$\eta$  is a model parameter

$\tilde{\alpha}(s)$  is the spatially-varying component of intercept  $\alpha$  at location  $s$

$\tilde{\beta}(s)$  is the spatially-varying component of parameter  $\beta$  at location  $s$

$\tilde{\eta}(s)$  is the spatially-varying component of parameter  $\eta$  at location  $s$

The spatially varying parameters ( $\tilde{\alpha}(s)$ ,  $\tilde{\beta}(s)$  and  $\tilde{\eta}(s)$ ) are assumed to follow mean-zero Gaussian processes ([Gelfand and Schliep, 2016](#)). A covariance function, defining covariance between two points to decay near-exponentially as the distance between them increases, is used to dictate the spatial behavior of the processes (see [Hunka et al. \(2025\)](#) for further details). Note, the observed values  $y(s)$  are often transformed for model fitting (e.g., square-root or cube-root transformation is applied to the NFI-estimated AGBD). The purpose of applying such

a transformation is that, first, it linearizes the relation between the observations and covariates (we assume such a linear relationship in eq. 1). Second, the transformation ensures the residuals (around a regression line) are approximately normally distributed and account for heteroskedasticity. In the case of Sudan, a cube-root transformation was found suitable. Upon model fitting, parameters  $\alpha$ ,  $\beta$ ,  $\eta$ ,  $\tilde{\alpha}(s)$ ,  $\tilde{\beta}(s)$  and  $\tilde{\eta}(s)$  are outputted, thereby allowing the prediction of vegetation AGBD at any given location with the EO-based covariates.

Model inference was conducted through Bayesian methods. In Bayesian methods, predictions are expressed as posterior predictive distributions, i.e., probability distributions of the unknown quantity based on observed data and model assumptions (for example, the distribution of model parameters is illustrated in Figure 2a). To predict vegetation AGBD with the model, numerical methods are required. This means, samples may be drawn from the posterior distributions of model parameters/effects multiple times, and predictions of AGBD may be made with these samples multiple times. The resulting posterior predictions of vegetation AGBD can be used to produce interpretable summaries, such as expected mean values of AGBD, their standard deviations (SDs) or uncertainty, and their 95% credible intervals (CIs). Such numerical approximations to predicted mean values, SDs, and 95% CIs of vegetation AGBD are illustrated further in Figure 2b and c, and described in equations below.




Figure 11: Approach to Bayesian inference and posterior prediction of vegetation AGBD at either a single location or across an arbitrary area. Equation numbers are indicated on this illustration to relate the steps to the described methodology.

Let  $s_*$  represent a desired prediction location (such as an unvisited NFI plot), where the posterior

$$y_{(m)}(s_*) = \left( \alpha_{(m)} + \tilde{\alpha}_{(m)}(s_*) \right) + \left( \beta_{(m)} + \tilde{\beta}_{(m)}(s_*) \right) x_1(s_*) + \left( \eta_{(m)} + \tilde{\eta}_{(m)}(s_*) \right) x_2(s_*) + \epsilon_{(m)}(s_*). \quad (2)$$

distribution of vegetation biomass must be approximated. Following the methods in [Hunka et al. \(2025\)](#),  $m = 1, \dots, M$  random samples are drawn from the posterior distribution of model parameters/effects. Samples from the pertinent parameters/effects are then plugged into the model (eq. 1) to yield posterior samples for vegetation AGBD,  $y(s_*)$ , at the location  $s_*$ :

Properties of the posterior distribution, such as the expected mean value,  $\bar{y}(s_*)$ , and SD,  $\sigma(s_*)$ , of AGBD, can then be approximated with the corresponding sample quantities,

$$\bar{y}(\mathbf{s}_*) = \frac{1}{M} \sum_{m=1}^M y_{(m)}(\mathbf{s}_*), \quad (3)$$

$$\sigma(\mathbf{s}_*) = \sqrt{\frac{1}{M-1} \sum_{m=1}^M (y_{(m)}(\mathbf{s}_*) - \bar{y}(\mathbf{s}_*))^2} \quad (4)$$

Over any arbitrary geographic area larger than the size of a single NFI plot, inference of area-wide mean AGBD and its associated uncertainty involves analyses of posterior predictions at multiple locations within the area. Assume the desired area,  $A$ , can be partitioned into a grid of  $N$  hypothetical NFI plots at locations  $\mathbf{s}_{1,*}, \dots, \mathbf{s}_{N,*}$ . The area-wide AGBD,  $y_{(m)}(A)$ , is then

$$y_{(m)}(A) = \frac{1}{N} \sum_{i=1}^N y_{(m)}(\mathbf{s}_{i,*}). \quad (5)$$

As  $m = 1, \dots, M$  random samples are drawn from the posterior distribution of model parameters, the mean area-wide AGBD,  $\bar{y}(A)$ , and standard deviation,  $\sigma(A)$ , is:

$$\bar{y}(A) = \frac{1}{M} \sum_{m=1}^M \bar{y}_{(m)}(A), \quad (6)$$

$$\sigma(A) = \sqrt{\frac{1}{M-1} \sum_{m=1}^M (\bar{y}_{(m)}(A) - \bar{y}(A))^2}. \quad (7)$$

To gauge the accuracy of the posterior predictions of vegetation AGBD (in comparison to the NFI estimates), an R-squared metric and root mean squared error (RMSE) serve as heuristics and are reported. These are estimated as,

$$\tilde{R}^2 = 1 - \frac{\sum_{i=1}^{t_*} (\mathbb{E}[y(\mathbf{s}_{*,i})] - y(\mathbf{s}_{*,i}))^2}{\sum_{i=1}^{t_*} (y(\mathbf{s}_{*,i}) - \bar{y}_*)^2} \quad (8)$$

$$\widetilde{\text{RMSE}} = \sqrt{\frac{\sum_{i=1}^{t_*} (\mathbb{E}[y(\mathbf{s}_{*,i})] - y(\mathbf{s}_{*,i}))^2}{t_*}} \quad (9)$$

where  $s_{*,i} ; i \in \{1, \dots, t_*\}$  are the locations with NFI-estimated  $y(s_{*,i})$ ,  $\bar{y}_*$  is the mean of those observed values, and  $E[y(s_{*,i})]$  are posterior mean predictions at the training or testing locations.

Further, the coverage rate of the 95% credible intervals (CIs), i.e. the proportion of NFI-estimates lying within the 95% s of our model AGBD predictions, is reported. A coverage rate of close to 95% is the ideal scenario. Coverage rates below 95% will indicate that model uncertainties may be overly optimistic, while rates above 95% suggests that uncertainties could be overly conservative (see [May et al., 2023](#); [Hunka et al., 2025](#)).

### 8.1.2.2. Assessing model results and validation

Model validation is conducted primarily through leave-one-out cross validation during the process of model fitting in R-INLA. A cross-validated Probability Integral Transform (PIT) is generated, which assesses, for each observation, the probability that the posterior mean prediction will be less than or equal to the observed value ([David and Johnson, 1948](#)).

$$\text{PIT}_i = \text{Prob}(y_i^{\text{pred}} \leq y_i^{\text{obs}} | y_{-i}), \quad (10)$$

where  $y_i^{\text{pred}}$  is the prediction where  $y_i^{\text{obs}}$  is the observed value, and  $y_{-i}$  denotes the observations with the  $i$ th observation omitted. When the model accurately represents the observations, the PIT scores are expected to be uniformly distributed between 0 and 1. Further, we report the coverage rate of the 95% credible intervals (CIs) of left-out samples. In the ideal case, approximately 95% of cases, the left-out sample will have a PIT score between 0.025 and 0.975.


### 8.1.2.3. Open-source Code Availability

The model for Sudan was run in R using package R-INLA ([R Core Team, 2022](#); [Lindgren and Rue, 2015](#)), which executes efficient geostatistical model fitting and the prediction of vegetation AGBD ([Gómez Rubio, 2020](#), Chapter 7). The source code is publicly released and is available on the NASA Carbon Monitoring Systems [Biomass Harmonization](#) GitHub repository. Two codes are presented: [Model fitting](#) and [Model predictions](#). Upon review, the code may be published with a permanent DOI record for use for Sudan's vegetation assessments in the future.

## 8.1.3. Results

### 8.1.3.1. Model performance

The geostatistical model reveals a strong relation between the covariates (i.e., NASA GEDI biomass and the Forest Non-Forest Probability Map 2023) and the observations (i.e., NFI-estimated AGBD) across Sudan (results in Figure 3). The model explains approximately 72% of the variability in the NFI-estimates of AGBD respectively (Figure 4). Corresponding model RMSEs are low, and there is no strong






Figure 12: (a) Results of the geostatistical model for prediction of aboveground biomass (AGBD), including model parameters and effects. Histogram shows the Probability Integral Transform (PIT). (b) Visualizations of the model results, including posterior distributions of the model parameters. Source code can be viewed at the link [Model fitting](#).

evidence of systematic deviations (empirical bias) in model predictions, with difference in means less than 1 Mg/ha. An appropriate coverage rate of the 95% CIs is noted. Model parameters, including the fixed effects ( $\alpha$ ,  $\beta$  and  $\eta$ ) and the spatially-varying random effects ( $\tilde{\alpha}(s)$ ,  $\tilde{\beta}(s)$  and  $\tilde{\eta}(s)$ ) are shown in Figure 3. The leave-one-out cross-validation resulted in a fairly uniform distribution of PITs (see Figure 3 and eq. 10), implying that the posterior samples of predicted AGBD do not have a larger probability of being over- or under-predictions when compared to their observed values. For the posterior predictions of AGBD, approximately 94.4% of PIT scores fall between values 0.025 and 0.975, indicating strong model performance.

### 8.1.3.2. Predictions of aboveground biomass

The primary purpose of using auxiliary EO datasets in support of the Sudanese NFI is allowing the prediction of AGBD in under-sampled areas due to accessibility and/or logistic constraints. Once the model parameters and associated spatial effects are estimated, samples are drawn from their posterior distributions. In a numerical approximation, predictions of AGBD can then be made with these posterior samples at any specific location (using eqs. 3 and 4) and across any desired area (using eqs. 6 and 7). A visualization of the prediction of AGBD at the locations of the whole NFI grid of Sudan is provided in Figure 5.




Figure 13: Posterior mean predictions of vegetation aboveground biomass density (AGBD) from the geostatistical model (eq 1), plotted against their NFI-estimated values. Source code can be viewed at the link [Model fitting](#).




Figure 14: Predictions of AGBD and their associated uncertainty, from the geostatistical model, at all locations of the Sudanese NFI. Source code can be viewed at the link [Model predictions](#).

Finally, for the estimation of emission factors per state, a conservative approach was adopted. The plots of the whole NFI grid were visually interpreted to identify whether the underlying land cover was “stable forest”, “forest degradation” or “forest gain”. Only when these conditions were met, posterior predictions of AGBD with the model parameters were made. State-wise summary statistics are provided in Table 14. Figures 1a-1m in **Annex 7** show a visual of the “stable forest”, “forest degradation” or “forest gain” locations of the NFI, and provide the model-predicted AGBD value at each of those locations.

**Table 14: Average above-ground biomass density by state**

State	AGBD [Mg/ha]	90% confidence interval [± %]
Blue Nile	37.94	50%
Central Darfour	13.89	25%
East Darfour	21	17%
El Gadarif	3.91	115%
Kassala	1.31	203%
North Darfour	4.26	37%
North Kordofan	7.76	102%
Sinnar	17.04	42%
South Darfour	22.57	23%
South Kordofan	16.91	21%
West Darfour	5.7	48%
West Kordofan	20.14	18%
White Nile	1.55	142%

Table 1: State-level predictions of vegetation aboveground biomass density (AGBD) estimates, and their uncertainty (standard deviation, SD90% confidence interval) across Sudan. Only GEDI predictions on NFI locations which are visually interpreted as stable forest, degraded forest or forest gain were included in the calculations.

## 8.2. Dead wood:

As explained in section 5.2 above, only standing deadwood has been included in this submission. The data on the volume of standing deadwood per state shown in Table 15 below has been obtained from the NFI (2021), converted to biomass using the weighted average wood density data in Table 16 below. The resulting values of biomass stock of the standing deadwood has been added to the living biomass stock of each state to obtained the emission factors for deforestation presented in Table 17 below.

**Table 15: Average aboveground volume/hectare (v/ha) of Standing deadwood disaggregated by state**

States	Standing Deadwood (m <sup>3</sup> /ha)	Sampling error (%)
North Darfour	0.2	155.4
South Darfour	5.0	61.3
West Darfour	1.9	147.3
East Darfour	2.2	67.3
Central Darfour	2.0	81.3
North Kordofan	0.8	114.0
South Kordofan	2.6	47.7
West Kordofan	2.9	29.1
Blue Nile	0.9	140.8
Sinnar	1.2	121.7
El-Gadarif	0.0	136.2

Source: NFI (2021) data

## 8.3. Emission Factors for deforestation:

Emission factors for deforestation have been estimated following the 2006 IPCC Guidelines, in particular equation 2.10 of Chapter two, using the average aboveground biomass density (V/ha) presented in Table 14 and dead wood data in Table 15 above. Default data for root shoot ratio value (0.332) of the tropical dry African region was obtained from the 2019 Refinement IPCC Guidelines table 4.4. and the carbon fraction (0.47) obtained from the 2006 IPCC Guidelines table 4.3. Wood density data for converting standing deadwood volume to biomass has been obtained from various sources including country specific data from national research, FAO, African Wood density Database of World Agroforestry Centre, Ethiopia FRL submission (2017) in addition to

the IPCC 2006 guidelines (see Annex 2 of this submission). The wood density values obtained from these sources are representative of RoS's forest circumstances. Weighted average wood density values have been calculated for each state based on the main tree species that represent more than 80% of the total wood volume in State (see Table 16 below).

**Table 16: Weighted Average Wood Density Values of the main Tree Species per State**

States	Weighted Average WD t. d.m/m <sup>3</sup>	No of main Species used in the calculation
North Darfour	<b>0.63</b>	19
South Darfour	<b>0.72</b>	18
West Darfour	<b>0.70</b>	18
East Darfour	<b>0.71</b>	18
Central Darfour	<b>0.69</b>	18
North Kordofan	<b>0.64</b>	19
South Kordofan	<b>0.73</b>	18
West Kordofan	<b>0.71</b>	18
Blue Nile	<b>0.64</b>	18
Sinnar	<b>0.69</b>	9
El-Gadarif	<b>0.64</b>	7

Emission Factors for each state has been calculated separately as the initial carbon stocks on Forestland converted to cropland, the results are presented in Table 17 below. The following equation 2.10 of the 2006 IPCC Guidelines have been used as the basis in the estimation of the initial carbon stock from living biomass and deadwood, as follow:

For living biomass using the equation:

$$G_{LB_j} = G_{W_j} * (1 + R) * CF * 44/12$$

Where:

$G_{BL}$  = average living (below- above ground) biomass stock on forestland converted to cropland, t.dm/ha

$G_W$  = average above-ground biomass stock, t.dm/ha (Table 14 above)

R = Root – shoot ratio (table 4.4, 19R IPCC Guidelines)

CF = Carbon fraction of dry matter, tonne C (t. d.m.)-1 (Table 4.3 2006 IPCC Guidelines)

44/12 = The ratio of molecular weights to convert C to CO<sub>2</sub>

j = State

For dead wood using the equation:

$$G_{DW_j} = I_{V_j} * WD_j * (1 + R) * CF * 44/12$$

Where:

$G_{DW}$  = average deadwood (below- above ground) biomass stock on forestland converted to cropland, t.dm/ha

$I_V$  = average above-ground volume of standing dead wood, m<sup>3</sup>/ha (Table 15 above)

$WD_i$  = weighted average wood density t. d. m /ha (Table 16 above)

R = Root – shoot ratio (table 4.4, 19R IPCC Guidelines)

CF = Carbon fraction of dry matter, tonne C (t. d.m.)-1 (Table 4.3 2006 IPCC Guidelines)

44/12 = The ratio of molecular weights to convert C to CO<sub>2</sub>

j = State

Based on the above estimates of biomass stock of living biomass and deadwood the total initial biomass carbon stock on forestland converted to cropland (EFs) have been calculated for each State as follow:

$$G_{Before_j} = (G_{LB_j} + G_{DW_j}) * CF * 44/12$$

Where:

$G_{Before}$  = average initial biomass carbon stock on forestland converted to cropland, t.CO<sub>2</sub>/ha

$G_{BL}$  = average living (below- above ground) biomass stock on forestland converted to cropland, t.dm/ha

$G_{DW}$  = average deadwood (below- above ground) biomass stock on forestland converted to cropland, t.dm/ha

CF = Carbon fraction of dry matter, tonne C (t. d.m.)-1 (Table 4.3 2006 IPCC Guidelines)

44/12 = The ratio of molecular weights to convert C to CO<sub>2</sub>

j = State

Table 17 below shows the emission factors ( $G_{Before_j}$ ) of the different States estimated based on the above explained methods, data and parameters. This is later used in equation 2.16 as the biomass carbon stocks on forest land before the conversion, see section 11.1 below. The 2006 IPCC Guidelines Approach 1 has been used to estimate uncertainty, in particular Equation 3.1 of

the 2006 IPCC Guidelines has been used to combine emission factor and other parameters used in the estimation of the EF for each State, with 90% confidence level.

**Table 17: Emission factors for deforestation per State**

States	Emission Factor (tCO <sub>2</sub> /ha)	Uncertainty 90% CI
Blue Nile	88.45	85%
Central Darfour	35.04	48%
East Darfour	51.83	39%
Gadarif	9.03	188%
Kassala	3.01	330%
North Darfour	10.10	65%
North Kordofan	19.03	158%
Sinnar	40.97	72%
South Darfour	60.10	46%
South Kordofan	43.19	44%
West Darfour	16.17	85%
West Kordofan	50.93	40%
White Nile	3.56	232%

## 9. Emission Factors for Forest Degradation:

As explained in section 5.1 above, Sudan lacks suitable data to develop emission factors to assess changes in carbon stock due to forest degradation with more accuracy. RoS does not have historical data and records from regular NFI measurements. The NFI (2021) report provide data based on measurements conducted during 2017-2020, however, is being established, with permanent sample plots to ensure regular measurements and updated data in future including changes of forest carbon stock over time and the effect of wood extraction and overgrazing, which are the main causes of forest degradation. Also, RoS lacks quality and regularly updated data on wood consumption, demand and supply, particularly of wood for energy.

Building on the exchange with Assessment Team during the technical assessment, Sudan decided to include forest degradation activity using emission factors based on average relative reduction in canopy cover over the reference period. In this context, canopy cover refers to the percentage of tree cover recorded during data collection. The relative reduction is the change in the canopy cover before and after forest degradation. During the reference data collection, Sudan gathered information on tree cover percentage under two specific conditions: first, when the land use and land cover (LULC) class was identified as "Forest," and second, across two distinct time periods: before and after a degradation event. The detailed response design used for data collection is outlined in Annex 9 of FRL, with the customized survey cards presented in Figures 4 and 5 of the Activity Data section.

The relative canopy cover reduction was calculating using the change in canopy cover observed before and after the degradation event, normalized by the canopy cover recorded in the last stable year before the observed degradation. As tree cover percentage data was available at the plot level, the relative difference was calculated at this scale and subsequently aggregated to the state level using the average function.

$$\text{Canopy Cover Reduction} = \frac{\text{Canopy Cover in Last Stable Year} - \text{Canopy Cover After Degradation}}{\text{Canopy Cover in Last Stable Year}}$$

Table 18 below shows the average relative reduction in the canopy over the reference period for each state where forest degradation occurs.

**Table 18: Average percentage reduction in canopy cover of forest degradation area per State**

Name of State	Average canopy Cover Reduction
Blue Nile	38%
Central Darfour	50%
Eastern Darfour	60%
El Gadarif	52%
Kassala	20%
Northern Darfour	50%
Sinnar	40%
Southern Darfour	40%
Southern Kordofan	32%
Western Darfour	58%
Western Kordofan	59%
White Nile	50%

For this submission, the emission factors for forest degradation are assumed to equal the changes in biomass carbon stock associated with the average relative reduction in the canopy cover in the forest degradation areas occurring during the reference period. The EFs for the deforestation (Table 17 above) are approximated as 100% biomass carbon stock on forestland before conversion is assumed to represent the initial biomass carbon stock loss on forestland, as post-deforestation carbon contents is equal to zero in Sudan, while the EFs for forest degradation in each state are approximated as the percentages loss in Table 18 multiplied with the biomass carbon stock on forest land (Table 17). Based on these assumptions, the EFs for forest degradation for each State have been calculated using the formula below. The emissions factors for forest degradation are presented in table 19 below.

$$EF_{FD_j} = \Delta Canopy_j * C_{STOCK_j}$$

Where:

**$EF_{FD}$**  = Emission factor for forest degradation per State, t.CO2/ha

**$\Delta Canopy$**  = Average relative (percentage) reduction in canopy cover of forest degradation (Table 18 above)

**$C_{STOCK_j}$**  = Average initial biomass carbon stock on forestland before degradation, t.CO2/ha (Table 17 above)

**j** = State

**Table 19: Emission factors for forest degradation per State**

States	Emission Factor (tCO <sub>2</sub> /ha)	Uncertainty 90% CI
Blue Nile	34.0	85%
Central Darfour	17.5	48%
East Darfour	30.9	39%
Gadarif	4.7	188%
Kassala	0.6	330%
North Darfour	5.1	65%
Sinnar	16.2	72%
South Darfour	23.9	46%
South Kordofan	13.8	44%
West Darfour	9.4	85%
West Kordofan	30.2	40%
White Nile	1.8	232%

## **10. The Activity Data for Enhancement of Forest Carbon Stock:**

The activity data (AD) for the estimation of the removals associated with the enhancement of forest carbon stocks due mainly to Afforestation and Reforestation (A/R) has been obtained from the Forest National Corporation (FNC) HQ and its offices in the States. FNC records are used instead of the spatially explicit gain data included in table 9, section 6 above, because the latter data includes different types of gains not only A/R as required for Sudan Enhancement FRL. It includes gains such as natural regeneration occurs on areas adjacent to forestland area, natural regeneration on agricultural schemes (croplands) left uncultivated (abandoned) for some years in addition to some of the annually planted A/R areas (young A/R forests are difficult to identify through remote sensing).

FNC has an official annual A/R programme which is being implemented on bare lands in degraded forest areas, in addition to afforestation activities on the agriculture schemes (cropland) in collaboration with farmers and local communities. FNC keeps good records of the annually planted A/R areas, based on annual reporting by FNC state-level offices and reports of supervision visits conducted to assess the implementation of these annual A/R activities. FNC records include all A/R areas planted as forests, including A/R planted by FNC and A/R planted by community and there is a separate column for each in the FNC annual reports, the latter is normally supported by FNC in terms of provision of seeds, seedlings and extension services. Most of the A/R community/private forests are registered according to their ownership, in line with provisions in the Forest Law. FNC also provides seedlings/seeds for tree planting on streets, villages, building, gardens, however, these do not meet the forest definition, FNC keeps records of the number of seedlings and the amount of seeds distributed to support these activities in their reports.

With the support of the FRL Taskforce (FRL TF) data of the annual A/R areas has been collected for the years of reference period (2012- 2021), disaggregated by states. The AD is also disaggregated by A/R implemented through seeds and seedlings. The AD does not include:

- A/R areas planted pre-2012, the starting year of the agreed reference period for this submission.
- Tree plantings on streets, villages, buildings, gardens which do not meet the definition of forest.
- Natural gains occurring as a result of natural regeneration in areas adjacent to forestland and on abandoned agriculture areas.

Data verification, quality checked of errors and missing records was performed using data from the State offices of FNC and through the consultation with the FRL TF members. Modifications have been made to the land area data to cover some of the missing records for West Kordofan State, which was established in year (2013) of the reference period to cover missing records of years 2012 and 2013. The modification was based on a simple assumption that the total annual A/R areas for year 2012 and 2013 planted by North and South Kordofan States are shared equally with the new state of West Kordofan, which is established on part of the land areas of North and South Kordofan States. Accordingly, a complete set of annual A/R area data has been obtained for all the 18 states of RoS for the years of the reference period (2012-2021), see Annex (1). In this data set, zero records in some years mean A/R activities have not been implemented.

Overall, the area planting through use of seeds represents about 90%, while the area planting through seedlings represents the remaining 20%, of the total area of the annual A/R activities in RoS. However, the success rate of the area planting by seeds is very low compared to the area planted by seedlings. This is generally attributed to lack of sufficient investment in the areas planted by seeds, including for land preparation, water harvesting, protection against grazing and other treatments such as weeding which is usually practiced once or twice to reduce weed competition. Planting by seeds in many cases is done simply through seeds broadcasting without any land preparation or other treatments. Higher success rates are to a large extend associated with the A/R areas supported through some donor funded projects, in which more investment is made including on nurseries, land preparation and other practices. Generally, data on success rate was difficult to obtain given the war circumstances in RoS and lack of access to the data in the buildings of FNC HQ and some state offices. In fact, FNC data does not conduct assessment of success rate of the annual A/R plantations for more than one or two years after their planting date. Accordingly, the available records may not be representative of the actual or average success rate of the A/R forests over the reference period. Therefore, estimates of the overall success rate of the annual A/R areas planted by seeds and seedlings has been obtained based on expert judgement, involving use of some of the data available in state offices, the expertise of the FNC staff working in the states and also in consultations with the members of the FRL TF. The estimates agreed on the average success rates of the A/R areas planted by seeds and seedling are presented in the table (20) below:

**Table 20: Success rates of A/R forests planted by Seeds and Seedlings for each State**

State	Success rate (Seeds) %	Entries used	Success rate (Seedlings) %	Entries used
<b>Northern</b>	25	<b>0.25</b>	68-70	<b>0.69</b>
<b>River Nile</b>	40-44	<b>0.42</b>	77	<b>0.77</b>
<b>Khartoum</b>	20-30	<b>0.25</b>	80	<b>0.80</b>
<b>Al-Gezira</b>	50	<b>0.50</b>	75	<b>0.75</b>
<b>Sinnar</b>	25-30	<b>0.28</b>	75	<b>0.75</b>
<b>White Nile</b>	45	<b>0.45</b>	60	<b>0.60</b>
<b>Blue Nile</b>	30-35	<b>0.33</b>	70-75	<b>0.73</b>
<b>Gadarif</b>	30-40	<b>0.35</b>	75	<b>0.75</b>
<b>Kassala</b>	35	<b>0.35</b>	70	<b>0.70</b>
<b>Red Sea</b>	20-30	<b>0.25</b>	70	<b>0.70</b>
<b>North Kordofan</b>	20-30	<b>0.35</b>	60	<b>0.60</b>
<b>South Kordofan</b>	50	<b>0.50</b>	53	<b>0.53</b>
<b>West Kordofan</b>	38	<b>0.38</b>	57	<b>0.57</b>
<b>North Darfour</b>	20-30	<b>0.25</b>	65-70	<b>0.68</b>
<b>South Darfour</b>	46	<b>0.46</b>	70	<b>0.70</b>
<b>East Darfour</b>	42-50	<b>0.46</b>	65-75	<b>0.70</b>
<b>West Darfour</b>	45-50	<b>0.48</b>	75-80	<b>0.78</b>
<b>Central Darfour</b>	35-45	<b>0.40</b>	70-80	<b>0.75</b>

A range of more than 12 multipurpose tree species are used by the different States of RoS in their annual A/R activities. These include tree species planted for nontimber forest products (NTFP) such as Gum production, other forest fruits, animal fodder and tree planted for round wood, timber, shade, etc. Different States use different combinations from these 12 tree species and plant them in different areas of their annual A/R programme area, this is in accordance with their local needs, climate, circumstances and objectives. Table 21 below include information on the names of the tree species used and the percentage of their areas from the total annual A/R area in the different States.

**Table 21: Common tree Species used in the annual A/R activities and percentages of their areas in each State**

States	Acacia senegal	Acacia seyal	Acacia nilotica	Acacia tortilis	Acacia mellifera	Khaya Senegale nsis	Eucalypt us Spp	Balanit es aegyptiaca	ziziphu s spp	Azadir achta indica	Dalber gia sissoo	Scleroc arya birrea
<b>Khartoum</b>			5	75			20					
<b>Al-Gezira</b>		10	30	35	20		5					
<b>Sinnar</b>	25	30	35		10							
<b>White Nile</b>	30	20	20	15	15							
<b>Blue Nile</b>	50	15	25		5	5						
<b>Gadarif</b>	10	62	10		10		3					
<b>Kassala</b>	42	45	2	4	2		5					
<b>Red Sea</b>				40	5			20	30	5		
<b>N. Kordofan</b>	62	5	5	10	10			4	4			
<b>S. Kordofan</b>	50	20	5		20							5
<b>W. Kordofan</b>	75	5	10		5			5				
<b>N. Darfour</b>	62		5	23	10					5	10	
<b>S. Darfour</b>	40				15	30						
<b>E. Darfour</b>	60	30			5	5						
<b>W. Darfour</b>	30		5		50					10		
<b>C. Darfour</b>	50		10		30	5			5			
<b>Northren</b>		15	8	40	2		35					
<b>River Nile</b>		20	10	42	10		15			3		

### **10.1. Removal Factors for Enhancement of Forest Carbon Stock:**

Data on annual growth rates of the species used in A/R is not available from national sources. The NFI (2021) report does not provide such data. Therefore, RoS used representative default Mean Annual Increment (MAI) values from table 4.11 of the 2019 Refinement IPCC Guidelines, which provides MAI values for all the native tree species used in RoS and this data is represents African conditions, except for *Azadirachta indica* MAI value, which was obtained from FAO data.

When selecting the MAI value for the A/R tree species, RoS selected MAI values of “Productive semi-natural forests”, because are representatives of the circumstances of its A/R forest formations that consist of productive native tree species, planted mostly by seeds on bare, degraded areas and generally take similar shape and provide the same function, as of the natural forest formation.

The wood density values of the main species used in A/R activities have been obtained from different sources including national research, the World Agroforestry Center, African Wood Density Database, etc, see Annex 2. The values of the root-shoot ratio are from 19R IPCC Guidelines table (4.4) and the Carbon Fraction from 2006 IPCC Guidelines table (4.3). Value of Biomass Expansion Factors (BEF) used in equation 2.10 was obtained from table 3A.1.10 of the IPCC GPG-LULUC (BEF1 values).

It is worth mentioning that in RoS's subnational FRL 2020, BEF values have not been used in the estimation of the A/R removals. However, in estimation of this FRL, it is noticed that the MAI data of table 4.11 of the 19R IPCC guidelines represents merchantable volume and not the total aboveground volume. Therefore, BEF1 value of the tropical broad leaf forest type has been used with wood density values to expand the merchantable volume to the total aboveground biomass. Table 22 below shows the parameters used in the estimation of the removal associated with forest enhancement activities in RoS.

**Table 22: Mean Annual Increment values and other parameters used in estimation of A/R removals**

Common species used in A/R activities	Growth rate 19R IPCC table 4.11		Wood Density	Biomass Expansion factor (BEF1)	Root shoot ratio		Carbon Fraction
	Mean Annual Increment (m <sup>3</sup> /ha/yr)**	St. Dev	t. dm/m <sup>3</sup>	Table 3A.1.10 GPG LULUCF	19R IPCC table 4.4	St. Dev	2006 IPCC table 4.3 tC/t. dm
<i>Acacia nilotica</i>	12.5-20	1.9	0.8	1.5 (1.3-1.7)	0.332	0.247	0.47
<i>Acacia seyal</i>	1.8-3.2	0.4	0.7	1.5 (1.3-1.7)	0.332	0.247	0.47
<i>Acacia senegal</i>	1.1-2.4	0.3	0.7	1.5 (1.3-1.7)	0.332	0.247	0.47
<i>Acacia mellifera</i>	1.9-3.5	0.4	0.7	1.5 (1.3-1.7)	0.332	0.247	0.47
<i>Acaica Tortilis</i>	1.2-3.7	0.6	0.44	1.5 (1.3-1.7)	0.332	0.247	0.47
<i>Khaya Senegalensis</i>	8.5-12	0.9	0.7	1.5 (1.3-1.7)	0.332	0.247	0.47
<i>Euclyptus Spp</i>	12-14	0.5	0.6	1.5 (1.3-1.7)	0.332	0.247	0.47
<i>Balanites aegyptiaca</i>	1.2-1.5	0.1	0.63	1.5 (1.3-1.7)	0.332	0.247	0.47
<i>ziziphus mauritiana</i>	0.9-1	0	0.6	1.5 (1.3-1.7)	0.332	0.247	0.47
<i>Sclerocarya birrea</i>	1.5-1.7	0.1	0.58	1.5 (1.3-1.7)	0.332	0.247	0.47
<i>Dalbergia sissoo</i>	4-6	3.8	0.6	1.5 (1.3-1.7)	0.332	0.247	0.47
<i>Azadirachta indica</i>	(4-19) 5 averg *	0.5	0.78	1.5 (1.3-1.7)	0.332	0.247	0.47

\*<https://www.fao.org/4/y9933e/Y9933E12.htm>

\*\*RoS used average values from the range in table 4.11

Based on these data sets, removal factors have been calculated for each state, representing a weighted average annual increment per hectare of the combination of the species and the

percentage of their areas in the annual A/R programme of each state. The calculation of the removal factors has been performed as follows:

Step 1: Use of equation 2.10 of the 2006 IPCC guidelines as the basis for the calculation of the annual increment of biomass carbon stock for each species used in A/R:

$$G_i = I_{V_i} * D_i * BEF_{I_i} * (1 + R_i) * CF * 44/12$$

Where:

$G_i$  = annual increment of biomass carbon stock of a specie used in A/R, tonne CO<sub>2</sub>/ha/year

$I_{V_i}$  = average net annual increment for specific vegetation type, m<sup>3</sup> ha/yr (Table 22 above)

$D$  = Wood density, tonnes d.m. m<sup>-3</sup> (Table 22 above)

$BEF_{I_i}$  = biomass expansion factor for conversion of annual net increment (including bark) to aboveground tree biomass increment, dimensionless; Table 3A.1.10 GPG-LULUCF

$R$  = ratio of below-ground biomass to above-ground biomass for a specific vegetation type, in tonne d.m. (19R IPCC Guidelines Table (4.4))

$i$  = Species

$CF$  = carbon fraction of dry matter, tonne C (tonne.d.m.)<sup>-1</sup>, 2006 IPCC Guidelines Table (4.3)

44/12 = The ratio of molecular weights to convert C to CO<sub>2</sub>

Step 2: estimate the weighted average annual increment of biomass carbon stock in A/R areas in each state, using the following formula:

$$G_{TOTAL_j} = \frac{\sum(G_i * Weight_i)}{total weight_{ji}}$$

Where:

$G_{TOTAL}$  = Weighted average annual increment of biomass carbon stock of all species used in A/R in each State, tonne CO<sub>2</sub>/ha/year

$G_i$  = annual increment of biomass carbon stock of a specie used in A/R, tonne CO<sub>2</sub>/ha/year

Weight = percentage of the area planted by a specie from the total areas the annual A/R in each State (Table 21 above)

Total weight = the Sum of weights of all species planted in the total A/R area in each State.

$j$  = State

$i$  = species

Uncertainty of the MAI and other parameters used in the calculation of the removal factors, has been propagated. However, FNC data on annual A/R areas do not include uncertainty values, therefore, it assumed zero. Hence, the uncertainty values here represent only uncertainty associated with the MAI and other parameters used in the estimation of the removal factors. Table (23) below contains the removal factors used in the estimation of the Enhancement FRL for each State.

**Table 23: Removal Factors, Weighted Average Annual Biomass Carbon Increment**

State	Weighted Average Annual Removal in A/R areas	
	(tCO <sub>2</sub> /ha/year)	Perc 90% CI
River Nile	12.32	18.3%
Northern	15.50	21.7%
Khartoum	10.39	21.9%
Red Sea	3.65	33.7%
Kassala	7.00	19.3%
El-Gadarif	10.30	21.0%
Blue Nile	15.76	25.8%
Sinnar	18.77	29.7%
Al-Gezira	17.97	27.7%
White Nile	12.96	25.1%
North Kordofan	6.37	20.8%
South Kordofan	7.35	18.1%
West Kordofan	8.41	23.8%
North Darfour	6.36	21.4%
South Darfour	12.12	22.6%
West Kordofan	8.21	24.5%
East Darfour	5.90	21.6%
Central Darfour	9.87	19.9%

## 11. APPROACHES AND METHODS APPLIED IN THE FRL CONSTRUCTION

The IPCC 2006 National GHGs Inventory Guidelines does not directly refers to the REDD+ activities. However, the UNFCCC decisions (4/CP.15 and 12/CP.17) require Parties to use the most recent IPCC Guidelines and to maintain consistency with their GHGs inventory in the estimation of FREL/FRL and REDD+ results. Accordingly, IPCC methodologies are widely used by all REDD countries in the construction of FREL/FRLs and estimation of the REDD+ results. RoS applies the 2006 IPCC Guidelines in the estimation of emissions/removals associated with the two REDD activities of deforestation and enhancement of forest carbon stock, taking into consideration its national circumstances.

### 11.1. Estimation of GHGs Emission from Deforestation:

Deforestation is calculated as the net average annual carbon emissions associated with forestland converted to non-forestland (mainly to cropland) over the reference period (2012- 2021), disaggregated by the States. RoS applies the 2006 IPCC guidelines equations 2.15 and 2.16 of Chapter 2, in the estimation of the emission associated with deforestation, as applicable in the circumstances of RoS.

The net average annual carbon emissions of deforestation is equal to the sum of the biomass carbon stock on the forestland areas converted to other land use in all States. Assuming the oxidation of all the biomass removed from these forest areas in the same year, because in RoS forests are mainly converted to annual cropping system, the removed wood is used for energy purposes, either directly as fuelwood or converted to charcoal, and is consumed in the same year. Accordingly, the estimation of net average annual emissions associated with deforestation follows the steps below:

**Step 1:** Obtain the annual area of deforestation, forestland converted to other lands (ATO\_OTHERS), to be used in equation 2.16 of the 2006 IPCC Guidelines. Data of annual deforestation (forest loss) has been obtained using remote sensing data as described in section 6 above.

**Step 2:** Calculate Initial biomass carbon stocks on Forestland converted to Cropland using equation 2.10. Table 17 above shows the Initial biomass carbon stock on forest land converted to cropland, which represent the carbon stock before the conversion ( $B_{Before_i}$ ) in equation 2.16 below.

**Step 3:** Calculate the carbon stocks change per area for the type of conversion (to cropland) applying equation 2.16 of the 2006 IPCC Guidelines. Values of carbon stocks in biomass immediately after conversion ( $B_{After_i}$ ) are assumed to be zero, since the land is cleared of all its forest vegetation before it is turned into annual crops cultivation.

Equation (2.16): Estimation of Annual Change in Biomass Carbon Stock on Land Converted to Another land Category (mainly to Cropland).

$$\Delta C_{Conversion_i} = \sum_i \{(B_{After_i} - B_{Before_i}) * \Delta A_{To-Others_i}\}$$

Where:

$\Delta C_{Conversion_i}$  = Change in biomass carbon stocks on land converted to another land category, tonnes CO<sub>2</sub>/Year (deforestation FREL per State)

$B_{After_i}$  = Biomass stocks on cropland immediately after the conversion, tonnes d.m. ha<sup>-1</sup>

$B_{Before_i}$  = biomass carbon stocks on forest land before the conversion, tonnes t.CO<sub>2</sub>/ha

$\Delta A_{To-Others}$  = area of land converted to another land-use category in certain year, ha/year  
 $i$  = State

**Step 4:** Estimate the deforestation FREL for each state using the 2006 IPCC Guidelines equation 2.15. As equation 2.15 is designed to be applicable for forestland conversion to all other land uses and in accordance with national circumstances. In the case of Sudan FREL, forest land is mainly converted to annual cropland and according to national circumstances the parameter  $\Delta C_G$  is assumed to equal zero because in the subsequent years after the conversion carbon gains in biomass from annual growth are offset by losses from their harvesting. Sudan has an annual cropping system in the rainfed sector of a of 3-6 months rotation, dominated by crops such as different varieties of sorghum and millet, sesame, sunflower, groundnuts. The parameter  $\Delta C_L$  is also assumed to equal zero as no losses due to wood harvesting, fuelwood removal or disturbances is expected on annual cropland. Accordingly, based on equation 2.15, deforestation FREL for each state will equals the value of the parameter  $\Delta C_{CONVERSION}$

$$\Delta C_B = \Delta C_G + \Delta C_{CONVERSION} - \Delta C_L$$

Where:

$\Delta C_B$  = Annual change in carbon stocks in biomass on land converted to other land-use category, in t.CO<sub>2</sub>/yr

$\Delta C_G$  = Annual increase in carbon stocks in biomass due to growth on land converted to another land-use category, in t.CO<sub>2</sub>/yr

$\Delta C_{CONVERSION}$ = Initial change in carbon stocks in biomass on land converted to other land-use category, in t.CO<sub>2</sub>/yr

$\Delta C_L$ = Annual decrease in biomass carbon stocks due to losses from harvesting, fuel wood gathering and disturbances on land converted to other land-use category, in t.CO<sub>2</sub>/yr

**Step 5:** Sum up the annual changes in biomass carbon stocks due to deforestation ( $\Delta C_{Conversion_i}$ ) of all the States to estimate the national deforestation FREL, the results are presented in table 24 below.

#### **11.1.1. Uncertainty:**

The 2006 IPCC Guidelines Approach 1 has been used to estimate uncertainty of the deforestation FREL for each State and for the national deforestation FREL, with 90% confidence level. As described in section 3.2.3.1 of volume 1 of 2006 IPCC Guidelines, Sudan estimates uncertainty by using the error propagation equation in two steps. First, the Equation 3.1 to combine emission factor, activity data and other parameters used in the estimation of deforestation FREL for each State. Second, the Equation 3.2 is used to estimate the overall uncertainty of national deforestation FREL. Table 24 below shows the uncertainty values associated with the values of the deforestation FREL for each State and the overall national FREL.

#### **11.2. Estimation of GHGs Emission from Degradation:**

Emissions associated with forest degradation are calculated as the average relative reduction in the biomass carbon stock before and after forest degradation occurs, during the reference period, in the areas identified as forest degradation in each state. The calculation of emissions from forest degradation can be summarized in the following steps:

**Step 1:** Obtain the annual area of forest degradation. Data of annual forest degradation per state has been obtained through remote sensing, the data is included Table 13 of section 6 above, which also provides the description of how the data has been developed.

**Step 2:** Obtain the emission factors for estimating emissions associated with forest degradation. The EFs equal the change in biomass carbon associated with average relative reduction in canopy cover in forest degradation area as calculated in section 9 and presented in Table 19 above.

**Step 3:** Calculate the carbon stock change in forest degradation areas in each State applying the following equation.

$$\Delta C_{FD_j} = EF_j + A_{FD_j}$$

Where:

$\Delta C_{FD_j}$  = Change in biomass carbon stock in forest degradation areas per State, t.CO2/yr

EF = Emission factors for forest degradation per State (Table 19 above)

$A_{FD_j}$  = Area of forest degradation per State, (Table 11 above)

j = State

**Step 4:** Sum the changes in biomass carbon stock ( $\Delta C_{FD_j}$ ) associated with forest degradation in all the States to obtain the national values of emission associated forest degradation during the reference period. The results are presented in Table 25 below

### 11.3. Estimation of Removal from Enhancement of Forest Carbon Stock:

Enhancement FRL is estimated as the average annual accumulated CO<sub>2</sub> removals from the A/R activities on cropland and bare land in degraded forest areas (mainly subjected to cultivation), over the reference period (2012-2021). RoS applies methods of the IPCC 2006 guideline for estimation of changes in biomass carbon stocks associated with A/R activities in these lands. In particular, RoS applies equations 2.10 and 2.9 as applicable in circumstances of RoS. The annual removals are estimated disaggregated by States and by A/R implemented by Seeds and Seedling separately.

**Step 1:** Obtain data on the area converted to Forest Land. The data on annual A/R areas was obtained from the records of the Forest National Corporation (FNC) HQ and State offices. The data is disaggregated by states and by areas planted by Seeds and areas planted by Seedlings. See Annex 1.

**Step 2:** Obtain data on the tree species used, their area percentage of the total annual A/R areas and the success rates for A/R forests by Seeds and A/R by seedling. This data represents the best estimates, based on some of the available FNC records (in states), expert observations and opinions from FNC staff at the different states, in addition to the consultation with the member of FRL TF. See Tables 20 and 21 above

**Step 3:** Using equation 2.10 of the 2006 IPCC Guidelines estimate the removal factor for each State. The removal Factor equal the average annual increment of biomass carbon stock estimated separately for A/R planted by Seeds and Seedlings, weighted by the area of each species used by the different States (see Table 23 above).

**Step 4:** Estimate the FRL for enhancement forest carbon stock for each state, which is calculated as the average annual increment of biomass stock accumulated in the annual A/R areas adjusted by the success rates of A/R forest in the different State, over the period 2012-2021. The enhancement FRL is estimated on the basis of Equations 2.9 of Chapter 2 of the 2006 IPCC Guidelines, as follow.

Equation 2.9: Average annual increment of biomass carbon stock in the annual A/R areas accumulated for the reference period.

$$\Delta C_{G_{STATE}} = \sum_{i,j} (A_{i,j} * G_{TOTAL} * Succ.\ rate_{i,j})$$

Where:

$\Delta C_{G_{STATE}}$  = Average annual accumulated increment of biomass carbon stocks on the annual A/R areas, tonnes CO<sub>2</sub>/Year

A = area of annual A/R planted by Seeds and Seedlings, ha

$G_{TOTAL_{i,j}}$  = average annual increment of biomass carbon stock in A/R area, tonnes CO<sub>2</sub>/ha/year (Table 23 above)

Succ. rate = percentage of the survival rate of the A/R plantation, (Table 20 above)

i = A/R area by seeds

j = A/R area by seedlings

STATE: Government subnational administrative level. RoS has 18 States

**Step 5:** Estimate the national FRL for Enhancement of Forest Carbon Stock as the sum of the averages annual increment of biomass carbon stocks accumulated in A/R area in all the States, as in the formula below, the results are presented in Table 26 below.

$$\textbf{Total CO2 removal} = \Delta C_{G_{STATE1}} + \dots + \dots + \Delta C_{G_{STATE18}}$$

Biomass loss due to wood removal or disturbance, in the A/R established forests is very unlikely to happen during the reference period, (trees will be 10 years old) and therefore, has not been including in the estimation of the annual removals. Wood harvesting in RoS happens only on forest plantations managed for specific rotation periods ranging, according to species, from 15 years to the maximum natural age of the tree species used in A/R. The specific objectives include protection of watershed, production of wood, timber and none wood forest product such as Gum Arabic, the latter is very common in most of the forest. Tree species such as Acacia Senegal and Acacia Seyal which are commonly used in most of the States, are mainly managed for production of Gum. These two species in addition to A. mellifera, A. tortilis and others

are also very important as animal fodder. Timber producing tree species such as *Acacia nilotica* and *Mahogani* are usually grown for more longer rotations, 25 year and above. Direct fuel wood collection by local people communities, is common in RoS, however, is mostly from deadwood in the natural forests and some mature plantation forests, therefore wood collection is not expected to happen in these 10 years old A/R forests.

Fires are important disturbance in RoS, however, according to NFI 2021, in most of the forest area in RoS there is no evidence of fire, only about 3.63% of the total forest area experienced burning. The most common type of fires is the surface fire, which represent about 92.1% of the burnt area, while crown fire represents only about 0.18% and the rest (about 7%) such as underground fire. According to the national fire management strategy framework (2021), fires in RoS are human induced (not natural fires) and are mostly surface fires, which burns surface litter, other loose debris and small vegetation and usually cause least damage to forests. The forest burnt areas are insignificant compared to the total burnt areas in country. This is might be attributed to high incidences of fire outbreak in dense grass area (savannah woodland) rather than forestland. Most of forest burn areas were found in Blue Nile, South Kordofan and South Darfur States. Fires incidences on A/R forest area are very unlikely because usually these are managed and better protected through fire lines. Regular data on fires monitoring is needed for the inclusion of possible emissions from fires in the national FREL/FRL, currently such data is not available.

## 12. RELEVANT POLICIES AND LEGISLATION

Forestry activities started in the RoS in 1901, when the Woods and Forests Ordinance was promulgated and the Department of Woodlands & Forests was established the same year. Adoption and implementation of administrative and legislative measures continued ever since. The first forest policy for the RoS was enacted in 1932 with a main objective to protect and reserve country forests resource. The policy included directives on afforestation and logging activities in and outside of reserved areas. Also, stated the prohibited actions in forests and outlined privileges and rights of the population in and surrounding the forests.

The Forest Act of 1932 divided the responsibilities of forest administration between the central and provincial authorities. In 1939 a royalty order was enacted for collecting royalties from forest products produced outside the forest reserves to discourage people from cutting trees outside the forest reserves.

In 1986, the revision of the 1932 policy was made to accommodate the constitutional, political, administrative, environmental and socio- economic changes. The revised policy of 1986 also recognized and encouraged establishment of private, community and institutional forests (including irrigated forest plantations in agricultural schemes). The policy stressed the role of people participation and forestry extension in the areas of forest plantation, management and protection. It provided clear directions to raise the total areas of natural forests, wildlife reserves, and range lands. Its main objectives were to reserve 20 percent of the total area of the country, manage the forest resources on sustainable basis, strengthening institutional capacity. In addition, to resolving inter-sectoral conflicts, which have been a major cause of deforestation.

The Forests Act 1989, since its promulgation, has been repeatedly praised as the most important piece of legislation in RoS. It legalized people's participation in forest management and recognized, under the FNC technical supervision, new types of forest ownership: private, community and institutional forest reserves to be managed by owners, committees and institutions respectively, in addition to the national and regional forest reserves.

The current Act of Forests and Renewable Natural Resources, 2002 (FRNR), provides the framework for integrated management and protection of forests and renewable natural resources encompassing pasture and range. The Act also obliges investors in agricultural schemes to leave not less than 10% of the total area of a rainfed project and not less than 5% of the area of an irrigated project as shelter belts and protective wind breaks.

The Forest National Corporation has been established by the Forest Act (1989). Among the missions of the FNC is to develop and implement public policies, rules, plans and methods for safeguarding and protection of the forest resources. FNC responsibilities include technical supervision of the forest resources, building awareness, promote afforestation and reforestation including development of the Gum Arabic and other forest products. FNC is currently responsible for the development and implementation of the national REDD+ programme.

## 13. PROPOSED FOREST REFERENCE LEVEL

### 13.1. Values of national FRL:

The RoS national Forest Reference Level includes the three REDD+ activities of reducing emissions from deforestation, reducing emission from forest degradation and enhancement of forest carbon stock. The reference period for three REDD+ activities is the same (2012-2021). The reference level values of the three REDD+ activities have been estimated disaggregated by the states of Sudan, as presented in tables 24, 25 and 26 below. Deforestation is identified in 13 states, over the reference period causes total net average annual emissions estimated at **9,317,358 t.CO<sub>2</sub>/year** with uncertainty value of 13% (Table 24). Forest degradation activity occurs in 12 states, over the reference period causes total net average annual emissions estimated at **1,678,433 t.CO<sub>2</sub>/year** (Table 25) with uncertainty value of 17%. Enhancement of forest carbon stock A/R activity covers all the 18 states of the country, over the reference period resulted in average annual accumulated removals estimated at **- 2,112,017 t.CO<sub>2</sub>/year** (Table 26), with uncertainty value of 7.34% however, this uncertainty estimates represent only the uncertainty of removal factors and related parameters as explained in section 10.1. and Table 23 above.

**Table 24: Proposed Values of the Deforestation FREL**

<b>States</b>	<b>Net Average Emissions from Deforestation</b>	
	<b>(t.CO<sub>2</sub>/Year)</b>	<b>CI 90%</b>
Blue Nile	904,979	33%
Central Darfour	185,653	56%
Eastern Darfour	676,226	28%
El-Gadarif	62,124	121%
Kassala	17,760	180%
Northern Darfour	33,382	61%
Northern Kordofan	176,780	77%
Sinnar	466,037	70%
Southern Darfour	1,960,391	44%
Southern Kordofan	2,875,668	20%
Western Darfour	34,882	183%
Western Kordofan	1,917,659	17%
White Nile	5,815	387%
<b>National Total</b>	<b>9,317,358</b>	<b>13%</b>

**Table 25: Proposed Values of the Degradation FREL**

<b>States</b>	<b>Average net annual emissions</b>	
	<b>(t.CO<sub>2</sub>/Year)</b>	<b>CI 90%</b>
Blue Nile	186,418	79%
Central Darfour	38,878	121%
Eastern Darfour	222,687	62%
El-Gadarif	10,057	166%
Kassala	182	369%
Northern Darfour	1,833	133%
Sinnar	64,333	69%
Southern Darfour	362,101	49%
Southern Kordofan	240,736	17%
Western Darfour	13,890	147%
Western Kordofan	536,134	9%
White Nile	1,184	165%
<b>National Total</b>	<b>1,678,433</b>	<b>17%</b>

**Table 26: Proposed Values of the Enhancement of Forest Carbon Stock FRL**

<b>State</b>	<b>Average Accumulated Annual Removal (t.C02/year)</b>
	<b>(t.C02/year)</b>
River Nile	60,490
Northern	41,491
Khartoum	201
Red Sea	7,156
Kassala	22,154
El-Gadarif	132,096
Blue Nile	41,554
Sinnar	112,758
Al-Gezira	97,699
White Nile	126,349
North Kordofan	417,151
South Kordofan	284,597
West Kordofan	393,040
North Darfour	49,143
South Darfour	151,072

West Darfour	86,344
East Drafour	46,267
Central Darfour	42,456
<b>National Total</b>	<b>- 2,112,017</b>

### **13.2. Description of change from Sudan's FRL 2020:**

RoS followed the stepwise approach, described in decision 12/CP.17, para II.10, to prepare this National scale Forest Reference Level (FRL). The preparation of the national FRL builds on the experiences, capacities, data and resources Sudan developed during the preparation of its subnational FRL (2020) as well as the feedback received through the UNFCCC technical assessment. In line with paragraph (e) of decision 13/CP.13, Table 27 below provides description of changes in Sudan's FRL 2025 as compared to Sudan's 2020 FRL submission.

**Table 27: Comparison of Sudan FRL 2020 and FRL 2025**

<b>FRL 2020</b>	<b>FRL 2025</b>
Scale: Subnational scale, representing 11% of the total forest land in Sudan, disaggregated by state (3 states)	Scale: National scale, covering all forest area and disaggregated by 18 states
Historical reference period 2006-2018	Historical reference period 2012-2021
REDD+ Activities:	REDD+ Activities:
<ul style="list-style-type: none"> <li>Reducing emission from deforestation</li> <li>Enhancement of forest carbon stock associated with A/R</li> </ul>	<ul style="list-style-type: none"> <li>Reducing emission from deforestation</li> <li>Reducing emissions from forest degradation</li> <li>Enhancement of forest carbon stock associated with A/R</li> </ul>
Pools: AGB and BGB	Pools: AGB, BGB and standing DW
Gases: CO <sub>2</sub>	Gases: CO <sub>2</sub>
AD methodology: Land cover and change maps for 4 change assessment points	AD methodology: updated methodology based on stratified area estimate approach
Deforestation FREL estimation:	Deforestation FREL estimation:
<ul style="list-style-type: none"> <li>EFs from NFI initial results,</li> <li>2006 IPCC Guidelines, equation 2.15 and 2.16 with default data for root shoot ratio and carbon fraction (section 4.5, Tables 4.3 and 4.4), overall average WD from 11 species.</li> </ul>	<ul style="list-style-type: none"> <li>Revised EFs based on NFI (2021) and GEDI data,</li> <li>2006 IPCC guidelines equations 2.10 and 2.16 with weighted average WD based on 7-19 main species for each state, R values from Table 4.4, 19R IPCC Guidelines, CF from Table 4.3 2006 Guidelines.</li> </ul>

<p>Enhancement FRL estimation:</p> <ul style="list-style-type: none"> <li>• Weighted average removal factors based on the species used in A/R in each state.</li> <li>• 2006 IPCC Guidelines equations 2.9 and 2.10</li> <li>• Values of the root-shoot ratio, the mean annual increment (MAI) and the Carbon Fraction was obtained from the 2006 IPCC Guidelines, section 4.5, Tables 4.4, 4.11B and 4.3 respectively. WD from national data.</li> </ul>	<p>Enhancement FRL estimation:</p> <ul style="list-style-type: none"> <li>• Weighted average removal factors based on the combination of species used in each state.</li> <li>• 2006 IPCC Guidelines equations 2.10 and 2.9</li> <li>• MAI data of Table 4.11 of the 19R IPCC guidelines, WD from national and regional sources, BEF values from Table 3A.1.10 IPCC GPG (2003), R values from 19R IPCC table 4.4 and CF values from 2006 IPCC Table 4.3</li> </ul>
---	--

## **14. ADJUSTMENT FOR NATIONAL CIRCUMSTANCES**

With splitting of Sudan in 2011 in two States, about half of the forest resources become part of the new state of South Sudan, while about two third of the population remains in RoS. This situation resulted in increasing pressure on the remaining forest resources and raised the need for improvements in policy, planning and management practices to respond to the new situation. The national REDD programme, which was initiated (2013) comes at an opportune time to review policy, institutional framework, assesses the forest resources and the related social and environmental frameworks. The processes and outcomes generated by the national REDD readiness programme enabled addressing most of the challenges and improvements needed to make the forest policy framework, planning and management practices relevant to addressing the circumstances created by the separation of South Sudan and responding to climate obligations. These include adoption of a National REDD Strategy, new forest definition, conduct of a NFI, development of MRV/NFMs and building of technical and institutional capacity within FNC and REDD related institutions at the national and subnational (state) level. The reference period selected for this FRL (2012-2021) considered representative of the current trend in terms of capturing the effect of all these national circumstances affecting forest sector. Accordingly, RoS does see the need to further undertake an adjustment to the proposed FRL in this submission. However, further work on the effects of policy development on forest management and implementation of REDD+ activities will be studied in the future updates of the national FREL/FRL.

## **15. UPDATING FREQUENCY**

The national FRL will be updated when there are relevant improvements in the methodologies, activity data and other parameters (e.g. NFI, remote sensing and national research). Development in RoS's climate obligations relevant to the forest and land use sectors and development in the international REDD+ framework are also important consideration in the assessment of needs for future updates of the FRL.

## **16. FUTURE IMPROVEMENTS**

Following the stepwise approach of REDD+ framework, RoS moved from the subnational FRL developed in 2020 to prepare a national FRL. The preparation of the national FRL has benefited from the knowledge, resources, data and experience developed during the preparation of the subnational FRL, including the feedback from the UNFCCC TA in 2020. However, RoS could not implement most of the findings on improvement provided by the UNFCCC TA. This is in part because there has been no significant change in the national circumstances regarding data during

the short time since the TA of the 2020 FRL and also because of the exceptional political and unrest situation during this period including the current war situation.

**Activity Data:** Resources permitted, FNC to integrate the Activity Data (AD) methodology used in the FRL submission into the National Forest Monitoring System (NFMS), to ensure that the system is regularly updated to reflect changes in forest activities. There is a need to strengthen the capacity of the RoS team in data collection and reporting processes, as an effort to improve both transparency and accuracy. There is a need for national definitions of all REDD+ activities be established in the RoS context to ensure uniform interpretation and monitoring moving forward. For instance, the definition of degradation used in this submission focuses primarily on reductions in canopy cover, but it may be useful to incorporate additional local factors to better capture the full scope of degradation.

**Emission Factors:** Future improvements of EFs of deforestation and forest degradation are mainly based on availability of good quality national data on wood volume and biomass density. The FNC has recently established MRV unit to improve data generation, documentation, reporting and data sharing. The unit needs to be strengthened and supported with adequate technical and institutional capacities to effectively performs its important responsibilities. The work of this unit is expected to improve data availability and quality on biomass stock, A/R, forest fires, emission factors, etc.

New and updated data on wood and biomass density will be generated through the regular measurements on the permanent sample plots established by the NFI (2021). The NFI (2021) report also includes a recommendation on improving data collection on soil carbon and dead wood pools. The EFs and other parameters will also be updated based on national research and studies planned as part of the national REDD+ programme. Currently Sudan is working with FAO on reviewing and improving the estimation of the total forest area and forest area by strata. This work is triggered by the discrepancies in the estimates of the forest area from number of sources, including the NFI (2021), Land Cover Map (2021). Based on this review an erratum is planned to be finalized, approved and annexed to the NFI report. In addition, the FNC is planning to seek support to conduct the second National NFI. This effort will build on the methodologies used in the 2021 NFI, while also addressing its associated shortcomings.

**Other REDD+ Activities:** Inclusion of other significant REDD+ activities namely degradation still remain an area for future improvement pending availability of the required data such as regular NFI measurements, high quality data on wood demand and consumption, etc. RoS included forest degradation in this submission after the exchange with Assessment Team during the technical

assessment, using emission factors based on average relative reduction in canopy cover over the reference period. The other REDD+ activities of conservation and sustainable management (SFM) of forests are currently not a priority for RoS, however, could be considered in future submission when good quality data become available for estimating emissions associated with their implementation.

**Deadwood:** The estimation of the amount of dead wood in RoS is difficult, because significant amount of deadwood is collected directly by local communities living in the proximity of the forests to meet their immediate energy demands. The Forest Product Demand Study (1995), estimated the amount of wood fuel directly collected by communities at about 72% of total fuelwood used in RoS and this is not captured in available records. The NFI (2021), provides estimates of carbon stock of deadwood including standing, fallen and stumps. These estimates are considered the first data on the carbon stock of dead wood remaining in the forests. However, RoS still does have the detailed data required by the 2006 IPCC Guidelines, such as data on the average annual transfer of carbon stock into and out of dead wood pools and data for the calculation of the difference between the dead wood carbon stock in two points in time. In this submission RoS includes only volume estimates of standing deadwood in the emission factor for deforestation, data on volume of standing deadwood considered by the NFI field experts as more reliable. Inclusion of dead wood remain an area for improvement in future updates when the require data become available. One of the recommendations in the NFI (2021) report states that future inventories should include data on forest soil carbon, litter and dead wood estimations.

**Inclusion of forest Fires:** The only important disturbance in RoS is fires, however, according to NFI 2021, in most of the forest area in RoS there is no evidence of fire, only about 3.63% of the total forest area experienced burning. The most common type of fires is the surface fire, which represent about 92.1% of the burnt area, while crown fire represents only about 0.18%) and the rest (about 7%) is underground fire. According to the national fire management strategy framework (2021), fires in RoS are human induced (not natural fires) and are mostly surface fires, which burns surface litter, other loose debris and small vegetation and usually cause least damage to forests. The forest burnt areas are insignificant compared to the total burnt areas in country. This might be attributed to high incidences of fire outbreak in dense grass area (savannah woodland) rather than forestland. Most of forest burn areas were found in Blue Nile, South Kordofan and South Darfur States. Also, most of the fires occurs on natural forest and very rarely on A/R and plantation forests, which are usually managed and better protected by fire lines. Good quality regular data on fires monitoring is needed for the inclusion of emissions from fires in the national FREL/FRL, currently such data is not available and therefore inclusion of emissions from forest fires remain and an area for improvement in future updates.

## 17. REFERENCES

Irrigated Forest Plantations in the Sudan: Case Study Gezira Agricultural Scheme  
By Talaat D. A. Magi <file:///Users/nagmeldin/Downloads/IrrigatedforestplantationintheSudan.pdf>

Alkaline pulping of some eucalypts from Sudan, P. Khristova <sup>a</sup>, O. Kordsachia <sup>b</sup>, R. Patt <sup>b</sup>, S. Dafaalla (2006): <https://www.sciencedirect.com/science/article/abs/pii/S0960852405002105>

Effect of Growth Rate on Wood Density of Eucalyptus Camaldulensis Wood of Coppice Origin Grown in White Nile State Sudan, Ma Malik A. Y. Abdelgadir: <https://www.semanticscholar.org/paper/Effect-of-Growth-Rate-on-Wood-Density-of-Eucalyptus-Malik-Abdelgadir/e86919792ed7bdb4f73aeeecd39afaf20e958dde8>

Sudan's Forest National Corporation (2021) National Forest Inventory Final Report:  
[https://www.fao.org/fileadmin/user\\_upload/faoweb/Themes\\_\\_pages/Forests/REDD-NFM/Sudan\\_MRV/Sudan - NFI\\_Report.pdf](https://www.fao.org/fileadmin/user_upload/faoweb/Themes__pages/Forests/REDD-NFM/Sudan_MRV/Sudan - NFI_Report.pdf)  
<https://openknowledge.fao.org/server/api/core/bitstreams/9d8fe4c5-b3c5-45c0-9cac-07c5b64d4c0c/content>

Sudan's REDD Programme (2018), In-depth Analysis of Drivers of Deforestation & Forest/Range Degradation

FNC (2021), Fire Management Strategy Framework

Sudan's Forest National Corporation (2021), The National REDD+ Strategy and Action Plan

GFOI (2020), Integration of remote-sensing and ground-based observations for estimation of emissions and removals of greenhouse gases in forests. Methods and Guidance from the Global Forest Observations Initiative

HCENR (2021), Sudan's updated Nationally Determined Contribution

Cochran, W. G. (1977) Sampling techniques. John Wiley & Sons.

FAO. 2016. Map Accuracy Assessment and Area Estimation: A Practical Guide. National forest monitoring assessment working paper No.46/E, 60p.

Kozak, M. (2011) Comparison of efficiency of geometric stratification and k-means algorithm in Univariate stratification of skewed populations. Int. J. Agricult. Stat. Sci., Vol. 7, No. 1, pp. 341-344.

Neyman, J. (1934) On the Two Different Aspects of the Representative Method: The Method of Stratified Sampling and the Method of Purposive Selection. Journal of the Royal Statistical Society, Vol. 97, 558-606.

Olofsson, P., Foody, G. M., Herold, M., Stehman, S. V., Woodcock, C. E., Wulder, M. A. 2014. Good practices for estimating area and assessing accuracy of land change. Remote Sensing of Environment, 148:42-57.

IPCC (2006) Guidelines for National Greenhouse Gases Inventory

IPCC (2019), Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories

FAO (2020) Global Forest Resources Assessment Report, Sudan

FNC (2020), Forest Reference Level Submission to the UNFCCC

Babcock, C., Finley, A.O., Andersen, H.E., Pattison, R., Cook, B.D., Morton, D.C., Alonso, M., Nelson, R., Gregoire, T., Ene, L., Gobakken, T., Næsset, E., 2018. Geostatistical estimation of forest biomass in interior Alaska combining Landsat- derived tree cover, sampled airborne lidar and field observations. *Remote Sensing of Environment* 212, 212–230. URL: <http://dx.doi.org/10.1016/j.rse.2018.04.044>, doi:10.1016/j.rse.2018.04.044.

Breidenbach, J., Ivanovs, J., Kangas, A., Nord-Larsen, T., Nilsson, M., Astrup, R., 2021. Improving living biomass C-stock loss estimates by combining optical satellite, airborne laser scanning, and NFI data. *Canadian Journal of Forest Research* 51, 1472–1485. URL: <http://dx.doi.org/10.1139/cjfr-2020-0518>, doi:10.1139/cjfr-2020-0518.

David, F.N., Johnson, N.L., 1948. The probability integral transformation when parameters are estimated from the sample. *Biometrika* 35, 182. URL: <http://dx.doi.org/10.2307/2332638>, doi:10.2307/2332638.

Dubayah, R., Blair, J.B., Goetz, S., Fatoyinbo, L., Hansen, M., Healey, S., Hofton, M., Hurt, G., Kellner, J., Luthcke, S., Armston, J., Tang, H., Dun- canson, L., Hancock, S., Jantz, P., Marselis, S., Patterson, P.L., Qi, W., Silva, C., 2020. The Global Ecosystem Dynamics Investigation: High-resolution laser ranging of the Earth's forests and topography. *Science of Remote Sensing* 1, 100002. URL: <https://www.sciencedirect.com/science/article/pii/S2666017220300018>, doi:<https://doi.org/10.1016/j.srs.2020.100002>.

Dubayah, R., Hofton, M., Blair, J., Armston, J., Tang, H., Luthcke, S., 2021. GEDI L2A Elevation and Height Metrics Data Global Footprint Level V002. URL: [https://lpdaac.usgs.gov/products/gedi02\\_av002/](https://lpdaac.usgs.gov/products/gedi02_av002/), doi:10.5067/GEDI/GEDI02\_A.002.

Duncanson, L., Kellner, J.R., Armston, J., Dubayah, R., Minor, D.M., Hancock, S., Healey, S.P., Patterson, P.L., Saarela, S., Marselis, S., Silva, C.E., Bruen- ing, J., Goetz, S.J., Tang, H., Hofton, M., Blair, B., Luthcke, S., Fatoyinbo, L., Abernethy, K., Alonso, A., Andersen, H.E., Aplin, P., Baker, T.R., Barbier, N., Bastin, J.F., Biber, P., Boeckx, P., Bogaert, J., Boschetti, L., Boucher, P.B., Boyd, D.S., Burslem, D.F., Calvo-Rodriguez, S., Chave, J., Chazdon, R.L., Clark, D.B., Clark, D.A., Cohen, W.B., Coomes, D.A., Corona, P., Cushman, K., Cutler, M.E., Dalling, J.W., Dalponte, M., Dash, J., de Miguel, S., Deng, S., Ellis, P.W., Erasmus, B., Fekety, P.A., Fernandez-Landa, A., Ferraz, A., Fischer, R., Fisher, A.G., García-Abril, A., Gobakken, T., Hacker, J.M., Heurich, M., Hill, R.A., Hopkinson, C., Huang, H., Hubbell, S.P., Hudak, A.T., Huth, A., Imbach, B., Jeffery, K.J., Katoh, M., Kearsley, E., Kenfack, D., Kljun, N., Knapp, N., Král, K., Krůček, M., Labrière, N., Lewis, S.L., Longo, M., Lucas, R.M., Main, R., Manzanera, J.A., Martínez, R.V., Mathieu, R., Memiaghe, H., Meyer, V., Men- doza, A.M., Monerris, A., Montesano, P., Morsdorf, F., Næsset, E., Naidoo, L., Nilus, R., O'Brien, M., Orwig, D.A., Papathanassiou, K., Parker, G., Philipson, C., Phillips, O.L., Pisek, J., Poulsen, J.R., Pretzsch, H., Rüdiger, C., Saatchi, S., Sanchez-Azofeifa, A., Sanchez-Lopez, N., Scholes, R., Silva, C.A., Simard, M., Skidmore, A., Stereńczak, K., Tanase, M., Torresan, C., Valbuena, R., Ver- beeck, H., Vrška, T., Wessels, K., White, J.C., White, L.J., Zahabu, E., Zgraggen, C., 2022. Aboveground biomass density models for NASA's Global Ecosystem Dynamics

Investigation (GEDI) lidar mission. *Remote Sensing of Environment* 270, 112845. URL: <https://www.sciencedirect.com/science/article/pii/S0034425721005654>, doi:<https://doi.org/10.1016/j.rse.2021.112845>.

Emick, E., Babcock, C., White, G.W., Hudak, A.T., Domke, G.M., Finley, A.O., 2023. An approach to estimating forest biomass while quantifying uncertainty and correcting bias in machine learning maps. *Remote Sensing of Environment* 295, 113678. doi:[10.1016/j.rse.2023.113678](https://doi.org/10.1016/j.rse.2023.113678).

Gelfand, A.E., Schliep, E.M., 2016. Spatial statistics and Gaussian processes: A beautiful marriage. *Spatial Statistics* 18, 86–104. URL: <https://doi.org/10.1016/j.spasta.2016.03.006>.

Gómez Rubio, V., 2020. Bayesian Inference with INLA. doi:[10.1201/9781315175584](https://doi.org/10.1201/9781315175584).

Hunka, N., May, P., Babcock, C., de la Rosa, J.A.A., de los Ángeles Soriano- Luna, M., Saucedo, R.M., Armston, J., Santoro, M., Suarez, D.R., Herold, M., Málaga, N., Healey, S.P., Kennedy, R.E., Hudak, A.T., Duncanson, L., 2025. A geostatistical approach to enhancing national forest biomass assessments with earth observation to aid climate policy needs. *Remote Sensing of Environment* 318, 114557. URL: <https://www.sciencedirect.com/science/article/pii/S0034425724005832>, doi:<https://doi.org/10.1016/j.rse.2024.114557>.

Lindgren, F., Rue, H., 2015. Bayesian Spatial Modelling with R-INLA. *Journal of Statistical Software* 63, 1–25. URL: <https://www.jstatsoft.org/index.php/jss/article/view/v063i19>, doi:[10.18637/jss.v063.i19](https://doi.org/10.18637/jss.v063.i19).

May, P., McConville, K.S., Moisen, G.G., Bruening, J., Dubayah, R., 2023. A spatially varying model for small area estimates of biomass density across the contiguous United States. *Remote Sensing of Environment* 286, 113420. URL: <https://www.sciencedirect.com/science/article/pii/S0034425722005260> doi:<https://doi.org/10.1016/j.rse.2022.113420>.

R Core Team, 2022. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. Vienna, Austria. URL: <https://www.R-project.org/>

## 18. ANNEXES

### Annex 1: Areas of the Afforestation and Reforestation of the 18 States of RoS

States	Northern		River Nile		Khartoum		Al-Gezira		Sinnar		White Nile	
Years	Seeds	Seedlings	Seeds	Seedlings	Seeds	Seedlings	Seeds	Seedlings	Seeds	Seedlings	Seeds	Seedlings
2012	748	5	1482	26	0	0	988	71	3802	105	1795	2683
2013	328	10	2990	39	3	1	1380	88	3297	423	68	81
2014	1134	19	1758	24	21	0	1723	42	1991	63	5371	128
2015	3066	42	2394	50	42	1	2318	4	4284	126	5932	186
2016	3677	34	2423	5	0	0	8132	15	6003	77	3738	1386
2017	2730	40	2423	5	0	0	0	0	4890	8	2415	0
2018	4626	14	794	0	0	0	643	0	2277	22	4049	12
2019	2310	13	852	0	25	0	0	150	4279	46	3048	67
2020	774	5	6	1785	0	0	654	42	215	0	4780	0
2021	363	12	2901	168	0	50	0	0	1927	383	5359	0

States	Blue Nile		El-Gadarif		Kassala		Red Sea		North Kordofan		South Kordofan	
Years	Seeds	Seedlings	Seeds	Seedlings	Seeds	Seedlings	Seeds	Seedlings	Seeds	Seedlings	Seeds	Seedlings
2012	258	5	5579	0	0	1316	1776	5	9735	579	9735	579
2013	1782	6	3726	8	525	273	865	5	7652	1826	7652	1826
2014	1971	5	3365	0	739	368	1791	5	36023	3986	12230	1856
2015	1470	0	2100	2100	672	294	1890	4	28728	6258	10752	2100
2016	980	5	9813	0	824	303	1680	0	85877	8666	11206	1698
2017	4205	1	13167	0	815	235	764	3	66734	1058	12961	1539
2018	1965	2	8307	0	1428	386	1448	2	96369	4434	20287	1940
2019	221	1	8274	0	1806	271	708	2	36189	0	24931	549
2020	164	3	4950	27	0	25	1134	0	55541	0	27260	1223
2021	229	0	10960	16	2058	420	630	1	93996	0	5262	1003

States	W. Kordofan		N. Darfour		S. Darfour		E. Darfour		W. Darfour		C. Darfour	
Years	Seeds	Seedlings	Seeds	Seedlings	Seeds	Seedlings	Seeds	Seedlings	Seeds	Seedlings	Seeds	Seedlings
2012	9735	579	1391	780	1733	394	382	265	762	913	630	46
2013	7652	1826	820	987	4000	346	777	336	2410	825	686	44
2014	8736	375	3488	1045	2294	212	1386	252	4683	335	941	168
2015	12306	1638	1512	1176	1680	252	8	42	1344	126	966	126
2016	17720	1197	4190	902	5675	323	5683	424	2872	35	903	240

<b>2017</b>	22680	1232	3700	888	10605	908	5880	588	1092	92	3045	258
<b>2018</b>	51054	2560	7136	1010	11804	376	12733	0	8679	135	2384	289
<b>2019</b>	55101	3262	6045	1629	3570	214	1680	2	7140	630	3276	297
<b>2020</b>	71192	3476	2401	1268	2662	861	1772	252	8190	449	1567	1884
<b>2021</b>	65252	7103	3140	1394	5040	294	4263	0	9240	890	4233	6182

## Annex 2: Wood Density data


Species	WD t. d.m/m <sup>3</sup>	Source
Acacia tortilis f. raddiana	0.44	FNC 2019, Integrated Carbon Sequestration Project Establishment of Biomass Carbon Baseline
Acacia Seyal	0.7	Tarig O. Khider and Osman T. Elsaki, 2012. Heat Value of Four Hardwood Species from Sudan, JOURNAL OF FOREST PRODUCTS & INDUSTRIES, 2012, 1(2), 5-9
Acacia Senegal	0.7	Tarig O. Khider and Osman T. Elsaki, 2012. Heat Value of Four Hardwood Species from Sudan, JOURNAL OF FOREST PRODUCTS & INDUSTRIES, 2012, 1(2), 5-9
Acacia Mellifera	0.7	Tarig O. Khider and Osman T. Elsaki, 2012. Heat Value of Four Hardwood Species from Sudan, JOURNAL OF FOREST PRODUCTS & INDUSTRIES, 2012, 1(2), 5-9
Acacia Nilotica	0.8	M. A. Elfldl, 1985. Biomass estimation and energy content of acacia nilotica in the Blue Nile Master thesis , University of Khratoum
Eucalyptus Spp	0.4 -0.7	Alkaline pulping of some eucalypts from Sudan, P. Khristova <sup>a</sup> , O. Kordsachia <sup>b</sup> , R. Patt <sup>b</sup> , S. Dafaalla (2006): <a href="https://www.sciencedirect.com/science/article/abs/pii/S0960852405002105">https://www.sciencedirect.com/science/article/abs/pii/S0960852405002105</a>
E. Camaldulensis	0.7	Effect of Growth Rate on Wood Density of Eucalyptus Camaldulensis Wood of Coppice Origin Grown in White Nile State Sudan, Ma Malik A. Y. Abdelgadir: <a href="https://www.semanticscholar.org/paper/Effect-of-Growth-Rate-on-Wood-Density-of-Eucalyptus-Malik-Abdelgadir/e86919792ed7bdb4f73aeeecd39afaf20e958dde8">https://www.semanticscholar.org/paper/Effect-of-Growth-Rate-on-Wood-Density-of-Eucalyptus-Malik-Abdelgadir/e86919792ed7bdb4f73aeeecd39afaf20e958dde8</a>
Khaya Senegalensis	0.6 - 0.85	Agroforestry Database 4.0 (Orwa et al.2009) <a href="https://apps.worldagroforestry.org/treedb/AFTPDFS/Khaya_senegalensis.PDF">https://apps.worldagroforestry.org/treedb/AFTPDFS/Khaya_senegalensis.PDF</a>
Dalbergia sissoo	0.62 -0.82	Agroforestry Database A tree reference and selection guide. Version 4.0. <a href="https://apps.worldagroforestry.org/treedb2/speciesprofile.php?Spid=645">https://apps.worldagroforestry.org/treedb2/speciesprofile.php?Spid=645</a>
Acacia polyacantha	0.81 – 0.9	World Agroforestry Center: African Wood Density Database Carsan S, Orwa C, Harwood C, Kindt R, Stroebel A, Neufeldt H, and Jamnadass R. 2012. African Wood Density Database. World Agroforestry Centre, Nairobi. <a href="https://apps.worldagroforestry.org/treesnmarkets/wood/data.php#">https://apps.worldagroforestry.org/treesnmarkets/wood/data.php#</a>
Acacia sieberana	0.65 -0.72	“World Agroforestry Center: African Wood Density Database
Albizia aylmeri	0.73 -0.8	“
Anogeissus leiocarpus	0.91 – 1.04	“
Azadirachta indica	0.65 – 0.9	“
Cupressus lusitanica	0.38 – 0.65	“
Dalbergia melanoxylon	1.23 – 1.33	“

Sclerocarya birrea	0.51-0.64	“
Boswellia papyrifera	0.720	Robert Nygård*and Björn Elfving (1999), Stem basic density and bark proportion of 45 woody species in young savanna coppice forests in Burkina Faso. <a href="https://hal.archives-ouvertes.fr/hal-00883170/document">https://hal.archives-ouvertes.fr/hal-00883170/document</a> .
Dalbergia melanoxylon	0.817	Robert Nygård*and Björn Elfving (1999), Stem basic density and bark proportion of 45 woody species in young savanna coppice forests in Burkina Faso. <a href="https://hal.archives-ouvertes.fr/hal-00883170/document">https://hal.archives-ouvertes.fr/hal-00883170/document</a> .
Albizia Amara	0.7	FAO: Appendix 1 - List of wood densities for tree species from tropical America, Africa, and Asia. <a href="http://www.fao.org/3/w4095e/w4095e0c.htm">http://www.fao.org/3/w4095e/w4095e0c.htm</a> Also in the IPCC 2006, chapter 4 table 4.13
Anogeissus leiocarpus	0.73	Ogunwusi, A.A. and Onwualu,A.P and 2Ogunsanwo, O.Y (2013) Comparative Analysis of Wood Properties of Afzelia africana and Anogeissus leiocarpus Growing in Nigeria. Chemistry and Materials Research <a href="http://www.iiste.org">www.iiste.org</a> ISSN 2224- 3224 (Print) ISSN 2225- 0956 (Online) Vol.3 No.3, 2013
Balanites aegyptiaca	0.63	IPCC 2006, Chapter 4, table 4.13
Albizia amara	0.70	IPCC 2006, Chapter 4, table 4.13
Tamarindus indica	0.81-1.14	<a href="https://apps.worldagroforestry.org/treensmarkets/wood/data.php#">https://apps.worldagroforestry.org/treensmarkets/wood/data.php#</a>
Acacia gerrardii var. gerrardii	0.9	<a href="https://prota.prota4u.org/protav8.asp?g=pe&amp;p=Acacia+gerrardii">https://prota.prota4u.org/protav8.asp?g=pe&amp;p=Acacia+gerrardii</a>
Boswellia serrata	0.5	<a href="https://www.fao.org/4/w4095e/w4095e0c.htm">https://www.fao.org/4/w4095e/w4095e0c.htm</a>
Terminalia macroptera	0.81-0.9	<a href="https://apps.worldagroforestry.org/treensmarkets/wood/data.php#">https://apps.worldagroforestry.org/treensmarkets/wood/data.php#</a>
Sterculia Spp	0.55	Gisel Reyes, Sandra Brown, Jonathan Chapman, and Ariel E. Lugo (1992) Wood Density of Tropical Tree Species
Adansonia digitata	0.3	<a href="https://apps.worldagroforestry.org/treedb/AFTPDFS/Adansonia_digitata.PDF">https://apps.worldagroforestry.org/treedb/AFTPDFS/Adansonia_digitata.PDF</a>
Terminalia spp	0.50, 0.51, 0.58 +	<a href="https://www.fao.org/4/w4095e/w4095e0c.htm">https://www.fao.org/4/w4095e/w4095e0c.htm</a>
Dalbergia melanoxylon	1.23 -1.33	<a href="https://apps.worldagroforestry.org/treensmarkets/wood/data.php#">https://apps.worldagroforestry.org/treensmarkets/wood/data.php#</a>
Combretum ghasalense	0.845	<a href="http://db.worldagroforestry.org//wd/species">http://db.worldagroforestry.org//wd/species</a>
Terminalia brownii	0.654	Ethiopia FRL submission to the UNFCCC : Average of genus ( <a href="http://db.worldagroforestry.org//wd/genus/Terminalia">http://db.worldagroforestry.org//wd/genus/Terminalia</a> )
Boswellia papyrifera	0.500	Ethiopia FRL submission to the UNFCCC
Sterculia setigera	0.320	Ethiopia FRL submission to the UNFCCC : <a href="http://db.worldagroforestry.org//wd/species/Schefflera_abyssinica">http://db.worldagroforestry.org//wd/species/Schefflera_abyssinica</a>
Bauhinia spp.	0.67	FAO Data: <a href="https://www.fao.org/4/w4095e/w4095e0c.htm">https://www.fao.org/4/w4095e/w4095e0c.htm</a>

Ficus sp.	0.32	“
Diospyros spp	0.82	“
Celtis spp.	0.59	“
Ziziphus mauritiana	535-1080 kg/m <sup>3</sup>	<a href="https://apps.worldagroforestry.org/treedb/AFTPDFS/Ziziphus_mauritiana.PDF">https://apps.worldagroforestry.org/treedb/AFTPDFS/Ziziphus_mauritiana.PDF</a>
Prosopis africana	0.91-1.04	<a href="https://apps.worldagroforestry.org/treesnmarkets/wood/data.php#">https://apps.worldagroforestry.org/treesnmarkets/wood/data.php#</a>
Terminalia macroptera	0.81- 0.90	“
Pseudocedrela kotschy	0.73- 0.80	“
Combretum aculeatum	0.474	Ethiopia FRL submission to the UNFCCC : <a href="http://db.worldagroforestry.org//wd/species/Scheflera_abyssinica">http://db.worldagroforestry.org//wd/species/Scheflera_abyssinica</a>
Commiphora africana	0.48	“
Ficus sycomorus	0.422- 0482	“
Acacia drepanolobium	0.769	“
Diospyros mespiliformis	0.85	<a href="https://apps.worldagroforestry.org/treedb/AFTPDFS/Diospyros_mespiliformis.PDF">https://apps.worldagroforestry.org/treedb/AFTPDFS/Diospyros_mespiliformis.PDF</a>
Prosopis africana	0.91-1.04	<a href="https://apps.worldagroforestry.org/treesnmarkets/wood/data.php#">https://apps.worldagroforestry.org/treesnmarkets/wood/data.php#</a>
Lannea fruticosa	0.515	Ethiopia FRL submission to the UNFCCC : <a href="http://db.worldagroforestry.org//wd/species/Scheflera_abyssinica">http://db.worldagroforestry.org//wd/species/Scheflera_abyssinica</a>
Stereospermum kunthianum	0.74	“
Pterocarpus lucens	0.7-0.8	<a href="https://apps.worldagroforestry.org/treesnmarkets/wood/data.php#">https://apps.worldagroforestry.org/treesnmarkets/wood/data.php#</a>
Lannea schimperi	0.515	Ethiopia FRL submission to the UNFCCC : <a href="http://db.worldagroforestry.org//wd/species/Scheflera_abyssinica">http://db.worldagroforestry.org//wd/species/Scheflera_abyssinica</a>
Commiphora Sp	0.389	“
Combretum glutinosum	0.9	GlobalWoodDensityDatabase.
Combretum hartmannianum	0.791	Average value for Combretum Sp: GlobalWoodDensityDatabase
Combretum gallabatense	“	“
Combretum paniculatum	“	“
Combretum lamprocarpum	“	“
Commiphora quadricincta	0.381	Average value for Commiphora Sp: GlobalWoodDensityDatabase
Capparis spinosa	0.691	Average value for Capparis Sp: GlobalWoodDensityDatabase
Guiera senegalensis	0.690	<a href="https://www.researchgate.net/publication/318379324_Variation_in_growth_wood_density_and_carbon_concentration_in_five_tree_and_shrub_species_in_Niger">John Charles Weber et.al (2018), Variation in growth, wood density and carbon concentration in five tree and shrub species in Niger: https://www.researchgate.net/publication/318379324_Variation_in_growth_wood_density_and_carbon_concentration_in_five_tree_and_shrub_species_in_Niger</a>

Acacia laeta	0.7	Average value of Acacia species in RoS
Lannea schweinfurthii	0.551	GlobalWoodDensityDatabase.
Maerua pseudopetalosa	0.58	Average of tropical Africa: Ethiopia FRL submission to the UNFCCC : <a href="http://db.worldagroforestry.org//wd/species/Scheflera_abyssinica">http://db.worldagroforestry.org//wd/species/Scheflera_abyssinica</a>
Albizia anthelmintica Albizia spp.	0.52	<a href="https://www.fao.org/4/w4095e/w4095e0c.htm">https://www.fao.org/4/w4095e/w4095e0c.htm</a>
Bauhinia rubescens	0.808	Average value for Bauhinia sp: GlobalWoodDensityDatabase.
Dobera glabra	0.67 - 0.71	<a href="#">World Agroforestry Centre   agroforestrytree database 4.0</a>
Cordia abyssinica Cordia Sp	0.53	<a href="https://www.fao.org/4/w4095e/w4095e0c.htm">https://www.fao.org/4/w4095e/w4095e0c.htm</a>
Acacia drepanolobium	0.7	Average value of Acacia species in Sudan
Mitragyna inermis	0.528	Average value for Mitragyna sp: GlobalWoodDensityDatabase.
Borassus aethiopum	1.02 -1.14	<a href="https://apps.worldagroforestry.org/treesnmarkets/wood/data.php#">https://apps.worldagroforestry.org/treesnmarkets/wood/data.php#</a>

### Annex 3: Sampling unit, plot and subplot design



## Annex 4: Calculation Procedures Used in the RoS NFI

### Introduction

This document details the main calculation procedures implemented in the National Forestry Inventory (NFI), especially regarding the allometric equations used to calculate derived variables such as volume, biomass, and carbon. The calculation procedures are stored as SQL instructions in the NFI database and can be executed using Silva Metricus software.

The calculation procedures described below are stored in the table named sibp\_calculationfields in the NFI database; this table is used by Silva Metricus software to perform calculations defined by the user and to store the results in the corresponding fields. The calculation procedures have a sequential order that allows using calculations based on other calculations; for example, to calculate volume per hectare, the number of trees per hectare must first be calculated to multiply this value by the volume of an individual tree. The table sibp\_calculationfields has the following fields:

Table : Fields of the table for storing descriptions of calculations

No.	Field	Type and length	Description
1	configurationnumber	Numeric	Configuration number of NFI
2	tablename	Short text	Name of the table containing the field to be calculated
3	fieldname	Short text	Name of the field to be calculated
4	presentationorder	Numeric	Calculation order
5	fieldname_p	Short text	Descriptive name of the field to be calculated in the primary language
6	fieldname_s	Short text	Descriptive name of the field to be calculated in the secondary language (not used in the case of the NFI of Sudan)
7	fielddescription_p	Short text	Broader description of the field to be calculated, which is done in the primary language.
8	fielddescription_s	Short text	Description or more extensive explanation of the field to be calculated, which is in the secondary language. (not used in the case of the NFI of Sudan)
9	fieldunits	Short text	Units of measurement for the corresponding calculation.
10	fieldtype	Short text	Type of field (text, number, list, etc.)
11	fieldlength	Numeric	Total length of the field
12	decimalsnumber	Numeric	Number of decimals to be used, in case it is a field of double precision.
13	fieldformat	Short text	Field display format; formed by the field length and the number of decimals.

No.	Field	Type and length	Description
14	useforcalculations	Logical	A field used to define whether the variable will be used in regression calculations (yes/no)
15	tablenameoldversion	Short text	Name of the table that contained the field in previous versions of SIBP2 (previous version of Silva Metricus)
16	fieldnameoldversion	Short text	Name of the field in previous versions of SIBP2.
17	calculationtype	Short text	Type of calculation: special or SQL; special is a type of calculation that is already predefined in Silva Metricus and SQL is a calculation that uses an expression in SQL language.
18	sqlexpression	Short text	SQL expression for the calculation.
19	specialexpression	Short text	Type of predefined calculation
20	specialtable	Short text	Special table associated with a predefined calculation
21	specialfield	Short text	Special field associated with the predefined calculation.
22	representdbh	Short text	Variable representing DBH in the predefined calculation.
23	representtotalheight	Short text	Variable representing total height in the predefined calculation.
24	representcommheight	Short text	Variable representing the commercial height or bole height in the predefined calculation.

Below, the calculations performed to estimate trees per hectare, basal area, volumes, biomass, and carbon for individual trees are described in detail.

## Tree-level Calculations

Below a description of the calculations for the tree table (aa\_tree) is detailed, most of the calculations make use of SQL instructions and are stored in the database so that they can be used in the variable calculation module of Silva Metricus.

### Trees per hectare

In the design of the forest inventory, sampling units of 4 rectangular plots of 20m x 250m are used in which all trees 20 centimeters or greater in DBH are measured; also, there are three nested plots (by SU) of 10m x 20m where trees with DBH between 10 and 19.9 centimeters are measured. Depending on the case, the trees have different hectare expansion factors, below are the expansion factors according to the DBH of the tree:

- $DBH \geq 10 \text{ cm}$  and  $DBH < 20 \text{ cm}$ : the total area of a plot where these trees are measured is  $200 \text{ m}^2$ , so in a sampling unit this type of trees is measured in an area of  $2400 \text{ m}^2$  ( $200 \text{ m}^2 \times 12$ ); therefore, the expansion factor is  $10000/2400 = 4.1667$ . This means that each tree of the indicated category represents 4.1667 trees per hectare.

- DBH≥20 cm: the total area of a plot where these trees are measured is 5000 m<sup>2</sup>, therefore, in a sampling unit this type of trees is measured in an area of 20000 m<sup>2</sup> (5000 m<sup>2</sup> x 4); in this case the expansion factor is 10000/20000=0.5. This means that each tree of the indicated category represents 0.5 trees per hectare.

At the database level, the number of trees per hectare that each of the measured trees represents is stored in the field tre\_treesperhectare of the table aa\_tree, it is a double number type of field, and the calculation corresponds to the following SQL instruction:

```
UPDATE aa_tree SET aa_tree.tre_treesperhectare = IIf([aa_tree].[tre_dbh]>=20, 0.5,
IIf([aa_tree].[tre_axisdistance]<=10 Or [aa_tree].[tre_axisdistance]>=240 Or
([aa_tree].[tre_axisdistance]>=120 And [aa_tree].[tre_axisdistance]<=130), 4.1667, 0.5))
```

### **Basal area**

The basal area per hectare represented by each of the trees measured in the field is calculated using the formula:

$$G = \left(\frac{DBH}{100}\right)^2 * 0.7854 * N$$

Where:

G=basal area in m<sup>2</sup>/ha

DBH=Chest Height Diameter or Reference Diameter in centimeters

N=Number of trees per hectare (calculated in the previous section)

The SQL instruction executed by Silva Metricus is as follows:

```
UPDATE aa_tree SET aa_tree.tre_basalarea = Round(([aa_tree].[tre_dbh]/100)^2*0.7854*
[aa_tree].[tre_treesperhectare],6)
```

Estimated total and commercial height

In the case of the NFI, it is not necessary to calculate the total and commercial height using regression models, as the heights are measured or estimated directly in the field for all trees.

### **Total and commercial tree volume**

For the calculation of the total volume<sup>3</sup> that each of the trees represents per hectare, a formula with a form factor of 0.6 is used and the volume of thick branches is added, the formula is as follows:

$$V_t = G * th * 0.6 + V_b$$

---

<sup>3</sup> It refers mainly to the volume of the trunk and large branches, excluding the stump.

Where:

$V_t$ =Volume of the tree in  $m^3/ha$

$G$ =basal area per hectare represented by the tree in  $m^2/ha$

$th$ =Total height of the tree in meters

$V_b$ = is the volume of the thickest branches, for the calculation the Smalian formula is used which is the average of the basal areas of the smaller and larger diameter multiplied by the length of the branch.

The field where the total bole volume in cubic meters per hectare is stored is `tre_totalvolumeha`, it is of double type and the calculation is made with 6 decimal places. The SQL instruction for the calculation is as follows:

### **Branch volume:**

```
UPDATE aa_tree SET aa_tree.tre_branchesvolumeha
=IIF(Round((IIF([aa_tree].[tre_branch1diameter]>0,([aa_tree].[tre_branch1diameter]/100)^2*0.7
854*[aa_tree].[tre_branch1length],0) +
IIF([aa_tree].[tre_branch2diameter]>0,([aa_tree].[tre_branch2diameter]/100)^2*
0.7854*[aa_tree].[tre_branch2length],0) +
IIF([aa_tree].[tre_branch3diameter]>0,([aa_tree].[tre_branch3diameter]/100)^2*
0.7854*[aa_tree].[tre_branch3length],0) +
IIF([aa_tree].[tre_branch4diameter]>0,([aa_tree].[tre_branch4diameter]/100)^2*
0.7854*[aa_tree].[tre_branch4length],0))*[aa_tree].[tre_treesperhectare],6)>0,Round((IIF([aa_tr
ee].[tre_branch1diameter]>0,([aa_tree].[tre_branch1diameter]/100)^2*0.7854*[aa_tree].[tre_bra
nch1length],0) +
IIF([aa_tree].[tre_branch2diameter]>0,([aa_tree].[tre_branch2diameter]/100)^2*0.7854*[aa_tree]
].[tre_branch2length],0) +
IIF([aa_tree].[tre_branch3diameter]>0,([aa_tree].[tre_branch3diameter]/100)^2*0.7854*[aa_tree]
].[tre_branch3length],0) +
IIF([aa_tree].[tre_branch4diameter]>0,([aa_tree].[tre_branch4diameter]/100)^2*0.7854*[aa_tree]
].[tre_branch4length],0))*[aa_tree].[tre_treesperhectare],6),0)
UPDATE aa_tree SET aa_tree.tre_volumeha =
Round(IIF(tre_boleheight>0,tre_basalareaha*tre_boleheight*0.56,
tre_basalareaha*(6.123*tre_dbh^0.442)*0.56),6)
```

### **Total volume:**

```
UPDATE aa_tree SET aa_tree.tre_totalvolumeha =
Round([aa_tree].[tre_basalareaha]*[aa_tree].[tre_totalheight]*[aa_tree].[tre_formfactortv] +
IIF([aa_tree].[tre_branchesvolumeha]>0,[aa_tree].[tre_branchesvolumeha],0),6)
```

For the calculation of the volume of the bole per hectare represented by each of the trees, the form factor 0.6 is always used; however, in this case the height of the bole is used not the total height, the SQL instruction is as follows:

```
UPDATE aa_tree SET aa_tree.tre_bolevolumeha =
Round([aa_tree].[tre_basalareaaha]*[aa_tree].[tre_boleheight]* [aa_tree].[tre_formfactorcv],6)
```

### Aboveground biomass

To estimate the aerial biomass (AGB) of the trees, the following biomass model is used:

AGB=0.0673\*(ρ\*DBH<sup>2</sup>\*th)<sup>0.976</sup>/1000, global equation developed by J. Chave et al.

Where:

AGB = aboveground biomass in megagrams per hectare [Mg/ha])

DBH = Diameter at Breast Height or reference diameter (cm)

th= total height in meters

ρ = wood density (g/cm<sup>3</sup>)

To use this allometric equation of aboveground biomass, the value of the wood density for each species in the database is required, so for some species (example *Acacia mellifera*) the average wood density that was reported in national studies was used (associated in the database). For establishing the wood density of other species in the database, the DRYAD<sup>4</sup> database was used as a reference. From this database, only species from the Africa region were used. If the species does not have a reference for average density in the DRYAD database, the average of the genus is used; if there are no genus data, the average of the family is used; if none of the above is possible to obtain, the average of all species is used, corresponding to 0.6035 g/cm<sup>3</sup>.

Before calculating the aboveground biomass, it is necessary to associate each of the trees with the wood density according to the species (field tre\_wooddensity), said procedure is done using the following SQL instruction:

```
UPDATE cs_treespecies INNER JOIN aa_tree ON cs_treespecies.tsp_code = aa_tree.tre_specie
SET aa_tree.tre_wooddensity = Round([cs_treespecies].[tsp_wooddensity],4)
```

For calculating the aboveground biomass in tons per hectare, the field tre\_biomassha is used and the SQL instruction is as follows:

```
UPDATE aa_tree SET aa_tree.tre_biomassha = Round((0.0673*([aa_tree].[tre_wooddensity]*
[aa_tree].[tre_dbh]^2*[aa_tree].[tre_totalheight])^0.976)/1000*[aa_tree].[tre_treesperhectare],6)
```

### Underground Biomass

The underground biomass (roots) of trees is stored in the field tre\_undergroundbiomass, which is a double precision field and uses megagrams per hectare as its units. The calculation of

---

<sup>4</sup> <http://datadryad.org/repo/handle/10255/dryad.235>

underground biomass uses the estimation of aboveground biomass as an independent variable, and the model by Cairns et al. (1997)<sup>5</sup> is applied for its calculation.

$$bgb = e^{[-1.0587 + 0.8836 \cdot \ln(agb)]}$$

Where:

bgb = underground biomass (Mg/ha)

agb = aboveground biomass (Mg/ha)

### **Carbon and CO<sub>2</sub> Equivalent**

Each species in the database is associated with a carbon percentage (47%), and this percentage is applied to each tree. The SQL statement for assigning the carbon percentage in the tree table is as follows:

```
UPDATE cs_treespecies INNER JOIN aa_tree ON cs_treespecies.tsp_code = aa_tree.tre_specie
SET aa_tree.tre_carbonpercentage =
IIF([cs_treespecies].[tsp_carbonpercentage]=Null,47,[cs_treespecies].[tsp_carbonpercentage])
```

To calculate the carbon amount, the biomass is multiplied by the carbon content (%). The amount of CO<sub>2</sub> equivalent is calculated by multiplying the carbon content by the constant 3.67. The following are the SQL instructions for calculating carbon and CO<sub>2</sub> equivalent:

#### **Aboveground Carbon:**

```
UPDATE aa_tree SET aa_tree.tre_carbonha = Round([aa_tree].[tre_biomassha]*
[aa_tree].[tre_carbonpercentage]/100,6)
```

#### **Underground Carbon (roots):**

```
UPDATE aa_tree SET aa_tree.tre_undergroundcarbon =
Round([aa_tree].[tre_undergroundbiomass]* [aa_tree].[tre_carbonpercentage]/100,6)
```

#### **Aboveground CO<sub>2</sub>:**

```
UPDATE aa_tree SET aa_tree.tre_co2ha = Round([aa_tree].[tre_biomassha]*
[aa_tree].[tre_carbonpercentage]/100*3.67,6)
```

#### **Underground CO<sub>2</sub> (roots):**

---

<sup>5</sup> Cairns, M.A., Brown, S., Helmer, E.H. & Baumgardner, G.A. (1997). *Root biomass allocation in the world's upland forests*. Oecologia, 111(1): 1–11.

```
UPDATE aa_tree SET aa_tree.tre_underground = Round([aa_tree].[tre_underground biomass]*[aa_tree].[tre_carbon percentage]/100*3.67,6)
```

### **Associated Land Use Class (LULC)**

To do estimates by type of LULC (at various levels) using the "ratio estimators" technique, it is necessary to associate each tree with the LULC in which it is located. The SQL instruction to assign the LULC point associated with each tree of the plot is as follows:

```
UPDATE aa_lucs INNER JOIN aa_tree ON (aa_lucs.luc_lucsnumber = aa_tree.tre_luccnumber)  
AND (aa_lucs.luc_plotnumber = aa_tree.tre_plotnumber) AND (aa_lucs.luc_tractnumber =  
aa_tree.tre_tractnumber) AND (aa_lucs.luc_measurementnumber =  
aa_tree.tre_measurementnumber) AND (aa_lucs.luc_inventorynumber =  
aa_tree.tre_inventorynumber) SET aa_tree.tre_lucscode = [aa_lucs].[luc_lucc]
```

### **LULC Code and Other Classification Levels**

Once the LULC number has been assigned to each tree, it is necessary to associate the corresponding LULC code, to perform calculations at the level of LULC and other land use or land cover classification systems. This assignment uses the table aa\_landusesubplot and the following SQL instruction:


```
UPDATE cc_landuselevel3 INNER JOIN aa_tree ON cc_landuselevel3.code =  
aa_tree.tre_lucscode SET aa_tree.tre_luclevel0 = [cc_landuselevel3].[luclevel0],  
aa_tree.tre_luclevel1 = [cc_landuselevel3].[luclevel1], aa_tree.tre_luclevel2 =  
[cc_landuselevel3].[luclevel2]
```

## Annex 5: List of Assets used for Activity Data

Sr. No.	Monitoring Year	Name of Asset	Asset Id
1	Time-2 (2023)	Landsat 2023	projects/ee-abdallatana/assets/LANDSAT_MOCIC_2023/RoS_FRE_L_Mosaic_landsat_2023_Project
2		Landsat 2024	projects/ee-moelmardi80/assets/Landsat8_Mosaic_RoS2024
3		Planet NICFI 2023	projects/ee-fardaadmad88/assets/RoS_FRE_Mosaic_landsat_2023_Project/RoS_planet_2023_30m
4		Planet NICFI 2024	projects/ee-fardaadmad88/assets/RoS_FRE_Mosaic_landsat_2023_Project/RoS_planet_2024_30m
5		ALOS 2023	projects/ee-nailayasmin-RoS/assets/ALOS_KandC2023_RoS
6	Time -1 (2011)	Landsat 2011	projects/ee-nailayasmin-RoS/assets/RoS_2011_2009_L457
7		ALOS 2010	projects/ee-nailayasmin-RoS/assets/ALOS_KandC2010_RoS
8	Training Data FNF (2010 – 2023)		projects/ee-nailayasmin-RoS/assets/ceo-Sdn_2020_LC_training_data-plot-data-2024-09-24_byjalal_edited
9	<b>RoS NFI Systematic grid (full population)</b>		projects/ee-nailayasmin-RoS/assets/RoS_NFI_grid_noduplicates

## Annex 6: Annual Composite of Landsat Year 2011


1. In order to create an annual composite of 2011, Optical mosaic recipe in SEPAL was used.
2. For AOI selection National customized boundary was used and uploaded as an asset in GEE, asset id is listed in Table-1.
3. Date range was extended beyond the 2011 as of insufficient data, as shown in screenshot below; season here defined as a full calendar year, so in this case date range used for composite is 2009-2011




4. The scene from only Landsat 4-5 are used to get a better quality composites for classification, comprises of both Tier-1 and Tier-2 collection



5. The pre-processing parameters are used are highlighted in below given screenshot



6. Missing pixels were filled using pixels from Landsat-7 collection



7. And finally, image composite was exported as an asset

## Annex 7: A visual of the “stable forest”, “forest degradation” or “forest gain” locations of the NFI

Figures 1a-1m below visualize the NFI locations identified as “stable forest”, “forest degradation” or “forest gain” per state, and provide the associated model prediction of AGBD with uncertainty at those locations.




Figure 1a: Predictions of AGBD and their associated uncertainty, from the geostatistical model, at locations of the Sudanese NFI identified as “stable forest”, “forest degradation” or “forest gain”.




Figure 1b: (...continued...)




Figure 1c: (...continued...)




Figure 1d: (...continued...)




Figure 1e: (...continued...)




Figure 1f: (...continued...)




Figure 1g: (...continued...)




Figure 1h: (...continued...)




Figure 1i: (...continued...)




Figure 1j: (...continued...)




Figure 1k: (...continued...)




Figure 11: (...continued...)




Figure 1m: (...continued...)

## Annex 8: Response Design

### Response Design and Explanatory variables for Activity Data IPCC land cover classes<sup>[11]</sup> Republic of Sudan, FLR, 2024

#### 1. Data Collection on LULC Classes for 2023

##### 1.1. Forestland

This category includes all land with woody vegetation consistent with thresholds used to define forest land in the national GHG inventory, sub-divided into managed and unmanaged, and by ecosystem type as specified in the IPCC Guidelines<sup>3</sup>. It also includes systems with vegetation that currently fall below, but are expected to exceed, the threshold of the forest land category.

Republic of Sudan's National Forest Corporation (FNC) defines forest as;  
"an area of land spanning at least a minimum area of 0.4 ha with trees that have attained or have the potential to attain at least 2 m in height and a minimum tree canopy cover of 10%. It includes windbreaks and/or shelterbelts with a minimum of 20 m in width".

##### Explanatory variables

- It does not include land that is predominantly under agricultural or urban land use.
- Forest is determined both by the presence of trees and the absence of other predominant land uses.
- The trees should be able to reach a minimum height of 2 meters in situ.
- Includes forest roads, firebreaks, and other small open areas; forest in national parks, nature reserves, and other protected areas such as those of specific environmental, scientific, historical, cultural, or spiritual interest.
- It **includes** the plantation for restoration purposes. Young natural stands and all plantations established for forestry purposes which have yet to reach a crown density of 10 percent or tree height of 2 m are included under forest, as are areas normally forming part of the forest area which are temporarily unstocked as a result of human intervention or natural causes, but which are expected to revert to forest." (FAO/UNEP, 1999)
- **Excludes** tree stands in **agricultural production systems**, such as fruit tree plantations, oil palm plantations, olive orchards, and agroforestry systems when crops are grown under tree cover.
- 

###### 1.1.1. Tree Cover Forest

Count the no. of yellow dots in plot falling over trees and record in the survey, each subplot represents 2 percent of plot.

## Explanatory Variable

- This option will only be available when LULC cover is forestland
- Tree cover forest refers to trees only in forest area
- Which means trees standing agricultural land or urban setting should not be included
- Respect the forest definition height when counting trees inside the forest area (2m)
- Do not confuse this with shrubland

## 1.2. Cropland

This category includes arable and tillage land, and agro-forestry systems where vegetation falls below the thresholds used for the forest land category, consistent with the selection of national definitions.

### Explanatory variables

- Includes arable and tillable land, rice fields, and agroforestry systems where the vegetation structure falls below the thresholds in the national definition used for the Forest land category (< 10% crown cover and < 20 m in width for windbreaks and/or shelterbelts).
- Cropland includes all annual and perennial crops.
- Annual crops include cereals, oils seeds, vegetables, root crops and forages.
- Perennial crops in combination with herbaceous crops (e.g., agroforestry) or as orchards, vineyards and plantations such as cocoa, coffee, tea, coconut, bananas
- Arable land, which is normally used for cultivation of annual crops, but which is temporarily used for forage crops or grazing as part of an annual crop-pasture rotation (mixed system) is included under cropland.
- Fellow land with and without trees

## 1.3. Grassland

This category includes **rangelands and pastureland** that is not considered as cropland. It also includes systems with vegetation that fall below the **threshold used in the forest land category and are not expected to exceed (trees < 2 meter in height)**, without human intervention, the threshold used in the forest land category. The category also includes **all grassland from wild lands to recreational areas** as well as **silvi-pastural systems**, subdivided into **managed and unmanaged** consistent with national definitions.

### Explanatory variables

- Grasslands can vary greatly in their degree and intensity of management, from extensively managed rangelands and savannahs – where animal stocking rates and fire regimes are the main management variables – to intensively managed (e.g. with fertilization, irrigation, species changes) continuous pasture and hay land.

- Grasslands generally have a vegetation dominated by **perennial grasses**, with **grazing as the predominant land use**, and are distinguished from “forest” by having a **tree canopy cover of less than 10 percent**.
- **Grasslands includes rangelands and pastureland that are not considered Cropland including systems with woody vegetation and other non-grass vegetation such as herbs and shrubs** (shrubs are trees where height is less than 2 meter).
- **Includes Savanna or savannah – mixed woodland-grassland ecosystem.**

## 1.4 Wetlands

This category includes land that is covered or **saturated by water for** all or part of the year (e.g., **peatland**) and that does not fall into the forest land, cropland, grassland or settlements categories. It includes reservoirs as a managed sub-division and natural rivers and lakes as unmanaged subdivisions.

### Explanatory variables

- Guidance is restricted to **Managed Wetlands where the water table is artificially changed** (e.g., drained or raised) or wetlands created through human activity (i.e., damming a river)
- Reservoirs or impoundments, for energy production e.g., Dam irrigation, navigation, or recreation (Flooded Land).
- **All water bodies**, including seasonal water bodies, swamps.
- **Wetlands Natural or artificial ponds.**
- **Rivers, Lakes and streams, waterfalls.**

## 1.5. Settlements

**This category includes all developed land, including transportation infrastructure and human settlements of any size, unless they are already included under other categories.**

### Explanatory variables

- Settlements include **residential, transportation, commercial, and production (commercial, manufacturing) infrastructure of any size**, unless it is already included under other land-use categories.
- The land-use category settlements include soils, herbaceous perennial vegetation such as turf grass and garden plants, trees in rural settlements, homestead gardens and urban areas.
- Examples of settlements include land along **streets, roads in residential (rural and urban) and commercial lawns, in public and private gardens, in golf courses** and athletic fields, e.g., cricket field and in parks, provided such land is functionally or administratively associated with **cities, villages or other settlement types** and is not accounted for in another land-use category.
- Airports, factories.

## 1.6. Other land

**This category includes bare soil, rock, ice, and all unmanaged land areas that do not fall into any of the other five categories. It allows the total of identified land areas to match the national area, where data are available.**

Explanatory variables

- Other Land includes **bare soil, rock, ice**, and all land areas that do not fall into any of the other five land-use categories.
- Other Land is often unmanaged.
- Active Mine dump generally but also include the dumps if not active.

## 1.7. Plantation

Includes commercial plantation like pine, eucalyptus, Acacia and others.

For Example, Citrus Trees near to riverbank in Northern, river Nile, Kassala and Khartoum states.



Another example is Date Palm tree near to riverbank in northern and river Nile state.



Date palm tree plantation

## 2. Recording Changes over the FRL Period (2023-2011)

### 2.1. Forest Loss (Deforestation)

Forest land changed to non-forest land during 2023-2011

This includes both changes in land use and reduction of canopy cover below forest threshold:

- When percentage tree cover is above 10% but land use change is observed as a major activity e.g. Forest land converted to cropland
- When percentage tree+ cover is reduced from  $>10\%$  to  $<10\%$  but no LULC change is observed
- Percentage tree cover reduced to  $< 10$  percent and LULC change is also observed
- Example: Heterogeneous plot: e.g. if a plot is showing 50% change but still has intact forest at 50%. Then considering context, if change is beyond the limit of plot that means its deforestation. If only inside the plot, then must flag that plot as no confidence for further review, should be flagged as “degradation”

**Tips:** Use the Majority rule while collecting data, always consider context before making decision to better understand the landscape, for example a plot might meet the forest definition criteria but if you zoom out and check the context it could be located under the urban settings or agricultural area, in that case this will not be considered as forest.

### **Examples**

- Forest land converted to plantation.
- Forest land converted to settlement.
- Forest land converted to cropland.
- 2018 was a forest and in 2022 converted to cropland and percentage tree cover is still meeting forest definition criteria (12 %), this will be classified as deforestation because Land use change is being observed
- 

### **2.2. Forest Enchantment (NF-F/ F- F)**

Forest enhancement is equivalent to the IPCC category of Land Converted to Forestland, but the conversion is being observed during the reference period (2011-2023)

**Forest enhancement in the context of RoS is defined as conversion of croplands and bare lands within degraded forest land use (mostly subject to cultivation) to forestland through human-induced afforestation and reforestation. This enhancement of carbon stock is being observed during reference period (2023- 2011).**

- Forest plantation/ regeneration on cropland being observed to exist first in or after year 2011 and continue to exist till 2023.
  - o Plantation over bare land near/within degraded forest land being observed to exist first in or after year 2011 and continue to exist till 2023).
- It's opposite of degradation.
- Should include all **afforestation and reforestation activities**.
- Should **not include natural succession**.

### **2.3. Forest Gain**

It should include **non-forest land converted to forest land**. But make sure it's meeting forest definition criteria (tree cover > 10% and 2m height).

- Its increase in tree canopy cover observed over reference period (e.g. from <10 % to >10%).
- It is opposite to forest loss, include all-**natural forest regeneration**.
- Natural succession of bare land degraded land.
- Not already addressed under enhancement class.

### **2.4. Forest Degradation (F-F)**

Forest degradation is defined as **reduction in canopy cover** in this assignment context.

- It will be a Forestland remains Forestland but only percentage tree cover reduced over time (2023- 2011).
- Reduction in canopy cover but total tree cover should remain **equal or greater than 10%**
- Examples** percentage of canopy cover is reduced to less than 30% in 2017 from 90% in 2016
- When canopy cover reduced to less than 10% even without LULC changes, that's **“Deforestation” NOT “Degradation”**.
- Reduction of canopy cover but still have percentage of forest cover as of forest definition.

## **2.5. NF – NF LULC Changes**

- Any LULC changes not addressed above will be included here.
- 

## **2.6. No Changes**

[11] [https://www.ipcc-nccc.iges.or.jp/public/gpglulucf/gpglulucf\\_files/Chp2/Chp2\\_Land\\_Areas.pdf](https://www.ipcc-nccc.iges.or.jp/public/gpglulucf/gpglulucf_files/Chp2/Chp2_Land_Areas.pdf)

## Annex 9: Members of Steering Committee, FRL Taskforce, Remote Sensing Team and Management and Technical Support Team

### Members of the National REDD+ Steering Committee

No.	Name	Gender	Institution
1	Anwar Abdelhameid	M	FNC/ Director General
2	Prof. Essam Warrag	M	FAO Representative
3	Mohamed Ahmed Elfadil	M	Ministry of Finance and economic development
4	Dr. Mona Mohamed Ali	F	Higher Council of Environment and Natural Resources
5	Adil Mohamed Ali	M	Sudanese Environmental Conservation Society
6	Representative		University of Khartoum
7	Dr. Nuha Mutwali	F	Forestry Research Centre
8	Abdelmonem Adris	M	Range and Pasture
9	Yahya Adam Abdalla	M	FNC
10	Sawsan Abdalla	F	FNC
11	Dr. Sayeda A. Khalil	F	REDD+ Coordinator

### Members of the FRL Taskforce

No.	Name	Gender	Institution/ State
1	Adam Mohamed Adam Abdullah	M	FNC/HQ
2	Amir Mohamed Ahmed Saleh	M	FNC/HQ
3	Mutasim Fadl Elseed Babiker	M	FNC/HQ
4	Manal Ali Yassin Ibrahim	F	FNC/HQ
5	Hanady Ibrahim Abdelgabbar Ali	F	FNC/HQ
6	Ali Haroon Ali Omer	M	University of Khartoum
7	Azza Ahmed Al-Tayeb	F	HCENR
8	Mohamed Awad Mohamed	M	Red See State/FNC
9	Hatim Nuh	M	RSSA
10	Safaa Ahmed Beraima Hamid	F	FNC/HQ
11	Mona Mohamed Rakhi	F	North Kordofan state/FNC
12	Fatim Abdelwahid Abdalla Khaleil	F	RSSA
13	Zahir Dafaalla Aboaglla Mohammed	M	El-Gadarif State/FNC
14	Osman Adam Babiker Adam	M	North State/FNC
15	Khalda Abbas Hassan Elgzouli	M	FNC/HQ
16	Osman Abdalla Osman Mohammed	M	FNC/HQ
17	Khidir Salah	M	REDD+ PMU

### Members of Republic of Sudan Remote Sensing Team Participated in Activity Data Collection

No.	Name	Gender	E-Mail	Institution
1	Mohamed Osama	M	osama.mohamed@fao.org	FAO

2	Fatim Abdelwahid Abdalla Khaleil	F	<a href="mailto:Abdallatana@gmail.com">Abdallatana@gmail.com</a>	RSSA
4	Wala Hassan Abdelrahman Musnad	F	walaa_musnad@yahoo.com	RSSA
5	Khalda Abbas Hassan Elgzouli	F	<a href="mailto:hassan_khalda@yahoo.com">hassan_khalda@yahoo.com</a>	FNC
6	Amani AmadEldin ElshiekhIdrees Mahmoud	F	emooemadaldin55@gmail.com	RSSA
7	Egbal Hashim Mohamed Elnageeb	F	<a href="mailto:egbal44@hotmail.com">egbal44@hotmail.com</a>	RSSA
8	Safaa Ahmed Beraima Hamid	F	<a href="mailto:safaaberaima2007@gmail.com">safaaberaima2007@gmail.com</a>	FNC
9	Solafa Babiker Mohamed Babiker	F	<a href="mailto:babiker.solafa@gmail.com">babiker.solafa@gmail.com</a>	RSSA
10	Hanady Ibrahim Abdelgabbar Ali	F	<a href="mailto:hanady552017@gmail.com">hanady552017@gmail.com</a>	FNC
11	Hanadi Kamal-eldin Ahmed Yagoub	F	<a href="mailto:hanadikamaleldin@gmail.com">hanadikamaleldin@gmail.com</a>	FNC
12	safa Khalid Yousif Alkhadir	F	safakhalid2016@gmail.com	RSSA
13	Omnia Omer Dffelseed Hamid	F	<a href="mailto:omniaomer4@gmail.com">omniaomer4@gmail.com</a>	RSSA
14	Mutasim Fadl Elseed Babiker	M	mutasim552017@gmail.com	FNC
15	Hatim Nuh	M	<a href="mailto:hatimoo22@yahoo.com">hatimoo22@yahoo.com</a>	RSSA
16	Hebatalla Eltayeb Ali Awadalla	F	Hebaeltayeb27@gmail.com	RSSA
17	Mohaned Eltijani Mohamed Elmardi	M	<a href="mailto:hody.mohamed@gmail.com">hody.mohamed@gmail.com</a>	RSSA
18	Manal Ali Yassin Ibrahim	F	<a href="mailto:manalfncorp76@gmail.com">manalfncorp76@gmail.com</a>	FNC
19	Zahir Dafaalla Aboaglla Mohmmed	M	zahrdfallh4@gmail.com	FNC

**Expert Team: Supporting National Forest Inventory with Earth Observation, a study conducted to prepared Revised Emission Factors for deforestation:**

No.	Name	Gender	Affiliation / address
1	Dr. Neha Hunka	F	University of Maryland/European Space Agency; ( <a href="mailto:neha.hunka@esa.int">neha.hunka@esa.int</a> ), <b>Contact person on this study</b>
2	Paul May	M	South Dakota School of Mines and Technology
3	Chad Babcock	M	University of Minnesota
4	Marieke Sandker	F	FAO Rome ( <a href="mailto:marieke.sandker@fao.org">marieke.sandker@fao.org</a> )
5	Naila Yasmin	F	FAO Rome ( <a href="mailto:naila.yasmin@fao.org">naila.yasmin@fao.org</a> )
6	Javier Garcia Perez (Gamarra)	M	FAO Rome ( <a href="mailto:javier.garciaperez@fao.org">javier.garciaperez@fao.org</a> )
7	Essam Warrag	M	Project Coordinator ( <a href="mailto:essamwarrag2@yahoo.com">essamwarrag2@yahoo.com</a> )
8	Osama Mohamed Eisa	M	Remote Sensing Expert ( <a href="mailto:osamahe4@gmail.com">osamahe4@gmail.com</a> )
9	Nagmeldin Mahmoud	M	FRL Expert ( <a href="mailto:goutbi@yahoo.com">goutbi@yahoo.com</a> )

**Management and technical support team**

No	Name	Gender	Role	Institution
1	Sayedah Ahmed Khalil	F	Coordinator, REDD+ PMU	FNC
2	Igbokwe Kennedy	M	Project Management support	FAO
3	Essam Warrag	M	Project Management support	FAO
4	Anwar Sidahmed	F	Project Management support	FAO

5	Marieke Sandker	F	Technical Support	FAO
6	Naila Yasmin	F	Technical Support	FAO
7	Osama Mohamed Eisa	M	Remote Sensing Expert	National consultant
8	Nagmeldin Mahmoud	M	FRL Expert	National consultant